



Mean-field interacting boson random point fields in weak harmonic traps

Hiroshi Tamura, Valentin A. Zagrebnov

► To cite this version:

Hiroshi Tamura, Valentin A. Zagrebnov. Mean-field interacting boson random point fields in weak harmonic traps. Journal of Mathematical Physics, 2009, 50 (2), 10.1063/1.3072911 . hal-00304344

HAL Id: hal-00304344

<https://hal.science/hal-00304344>

Submitted on 22 Jul 2008

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Draft: July 16, 2008

Mean-Field Interacting Boson Random Point Fields in Weak Harmonic Traps

Hiroshi Tamura¹

Graduate School of the Natural Science and Technology
Kanazawa University,
Kanazawa 920-1192, Japan

Valentin A.Zagrebnov²

Université de la Méditerranée(Aix-Marseille II) and
Centre de Physique Théorique - UMR 6207
CNRS-Luminy-Case 907, 13288 Marseille Cedex 9, France

Abstract

A model of the mean-field interacting boson gas trapped by a weak harmonic potential is considered by the *boson random point fields* methods. We prove that in the Weak Harmonic Trap (WHT) limit there are two phases distinguished by the boson condensation and by a different behaviour of the local particle density. For chemical potentials less than a certain critical value, the resulting Random Point Field (RPF) coincides with the usual boson RPF, which corresponds to a non-interacting (*ideal*) boson gas. For the chemical potentials greater than the critical value, the boson RPF describes a divergent (local) density, which is due to *localization* of the macroscopic number of condensed particles. Notice that it is this kind of transition that observed in experiments producing the Bose-Einstein Condensation in traps.

Key words: Boson Random Point Field, Weak Harmonic Trap, Non-Homogeneous Bose-Einstein Condensation, Mean-Field Interacting Bose-Gas.

¹tamurah@kenroku.kanazawa-u.ac.jp

²zagrebnov@cpt.univ-mrs.fr

Contents

1	Introduction and Main Results	2
1.1	Weak Harmonic Traps	2
1.2	Ideal and Mean-Field Boson Gases in the WHT	3
1.3	Main Results	7
2	Preliminary Arguments and Estimates	10
2.1	$\Xi_\kappa(\beta, \mu)$ and $\tilde{\Xi}_\kappa(\beta, \mu)$	10
2.2	Approximations of One-Particle Gibbs Semigroups	12
2.3	Estimates for the Scaled Mean-Field Interaction	16
2.4	Evidence of Two Thermodynamic Regimes	18
3	Proof of Theorem 1.2	20
3.1	The case $\mu \leq \mu_c(\beta)$ (normal phase).	20
3.2	The case $\mu > \mu_c(\beta)$ (condensed phase).	24
4	Proof of Theorem 1.8	29
5	Concluding Remarks and Conjectures	30
A	Miscellaneous formulae	32

1 Introduction and Main Results

1.1 Weak Harmonic Traps

We consider the quantum statistical mechanical models of boson gas equipped with a κ -parameterized family of one-particle Hamiltonians of harmonic oscillators:

$$h_\kappa = \frac{1}{2} \sum_{j=1}^d \left(-\frac{\partial^2}{\partial x_j^2} + \frac{x_j^2}{\kappa^2} - \frac{1}{\kappa} \right), \quad (1.1)$$

which are self-adjoint operators in the Hilbert space $\mathfrak{H} := L^2(\mathbb{R}^d)$.

In this setup a "thermodynamic limit" corresponds to $\kappa \rightarrow \infty$ (i.e. the "opening" of the trap [DGPS]), which we call the Weak Harmonic Trap (WHT) limit. Notice that the set $C_0^\infty(\mathbb{R}^d)$ is a form-core of the operator (1.1) and that this set is also a form-core for the operator $(-\Delta)/2$. Here Δ is the standard Laplace operator in \mathbb{R}^d . Then (see e.g. [Ka]) one obtains the strong resolvent convergence:

$$\lim_{\kappa \rightarrow \infty} h_\kappa = (-\Delta)/2. \quad (1.2)$$

In spite of convergence (1.2), there is a drastic difference between the properties of the infinite Ideal Boson Gas (IBG) systems prepared via the WHT limit and via standard *thermodynamic limit* (TDL) $\lim_{L \rightarrow \infty} \Lambda_L = \mathbb{R}^d$ with the one-particle operators $t_L := (-\Delta/2)_L$ with a "non-sticky" (e.g. Dirichlet) boundary conditions [VVZ]. Here $\Lambda_{L=1} \subset \mathbb{R}^d$ is a bounded open region of unit volume $|\Lambda_{L=1}| = 1$ containing the origin $x = 0$ whose boundary $\partial\Lambda_{L=1}$ is piecewise continuously differentiable and

$$\Lambda_L := \{x \in \mathbb{R}^d | L^{-1}x \in \Lambda_{L=1}\}, \quad L > 0. \quad (1.3)$$

In fact, it is known that the Bose-Einstein Condensation (BEC) occurs for dimensions $d > 1$ in the IBG via WHT limit $\kappa \rightarrow \infty$, while for dimensions $d > 2$ in the IBG via TDL, see (1.13) and [DGPS], [PeSm]. Similarly, it is well-known that thermodynamic properties of the boson gases are very sensible to different ways of taking the thermodynamic limit [LePu] or to the different choices of the boundary conditions [VVZ].

The purpose of the paper is to examine the position distribution of the mean field boson model in WHT limit and to compare its behaviour to those of the mean field boson models in TDL or ideal boson models in WHT limit.

Our method is based on the theory of Random Point Fields (RPFs) (see e.g. [DV]). The usual boson and the fermion RPFs [Ly, M75, M77] have been formulated in a unified way in terms of the *Fredholm determinant* together with other related RPFs, which are indexed by fractional numbers, in [ShTa]. They have been re-derived as theories which describe position distributions of the constituent particles of quantum gases in the thermodynamic limit for canonical ensembles in [Tia]. It was shown that the Random Point Field (RPF) corresponding to fractional numbers [ShTa], describes the gases which consist of particles obeying the parastatistics [Tic]. The RPF describing a homogeneous Bose-Einstein condensation have been studied for the first time in [Tib], where the RPF is given by the convolution of usual boson RPF and another RPF. The latter one seems to describe position distribution of the condensed part of the constituent bosons. This RPF has been re-formulated using the Cox process.[EK]

These theories of RPFs yield a precise information about the *position* distribution of the constituent quantum particles, although they are not suitable to characterize the quantum systems completely (however see [FF, F]).

1.2 Ideal and Mean-Field Boson Gases in the WHT

Recall that in the *grand-canonical* Gibbs ensemble the partition function of the IBG trapped by harmonic potential (1.1) is given by

$$\Xi_{0,\kappa}(\beta, \mu) := \sum_{n=0}^{\infty} e^{\beta\mu n} \operatorname{Tr}_{\mathfrak{H}_{symm}^n} [\otimes^n G_{\kappa}(\beta)]. \quad (1.4)$$

Here $\mathfrak{H}_{symm}^n := (\otimes^n L^2(\mathbb{R}^d))_{symm}$ is the n -fold *symmetric* Hilbert space tensor product of $\mathfrak{H} := L^2(\mathbb{R}^d)$, $G_{\kappa}(\beta) = e^{-\beta h_{\kappa}}$ the one particle Gibbs semigroup. The zeroth term in (1.4) equals to 1 by definition. We consider the case of positive inverse temperature $\beta > 0$ and of negative chemical potential $\mu < 0$.

The spectrum of the operator (1.1) is discrete and has the form:

$$\text{Spec}(h_\kappa) = \{ \epsilon_\kappa(s) := |s|_1/\kappa \mid s = (s_1, \dots, s_d) \in \mathbb{Z}_+^d \} \quad (1.5)$$

where $|s|_1 := \sum_{j=1}^d s_j$ and \mathbb{Z}_+^1 is the set of all non-negative integers. The normalized eigenfunctions of the states for $s \in \mathbb{Z}_+^d$ are given by

$$\phi_{s,\kappa}(x) = \frac{1}{\kappa^{d/4}} \phi_s(x/\sqrt{\kappa}) = \prod_{j=1}^d \frac{1}{\kappa^{1/4}} \phi_{s_j}(x_j/\sqrt{\kappa}) , \quad (1.6)$$

where for each component j , ϕ_{s_j} is related to the *Hermite* polynomials $H_{s_j}(z)$ by

$$\phi_{s_j}(x_j) = (2^{s_j} s_j! \sqrt{\pi})^{-1/2} H_{s_j}(x_j) e^{-x_j^2/2}, \quad j = 1, \dots, d . \quad (1.7)$$

The ground state is denoted in this paper by

$$\Omega_\kappa(x) = \frac{1}{(\pi\kappa)^{d/4}} e^{-|x|^2/2\kappa} \equiv \phi_{s=0,\kappa}(x) , \quad (1.8)$$

where $x = (x_1, \dots, x_d) \in \mathbb{R}^d$, $|x|^2 := \sum_{j=1}^d x_j^2$.

Integral kernel of $G_\kappa(\beta) = e^{-\beta h_\kappa}$ has the explicit form (the Mehler's formula for *oscillator processes*):

$$G_\kappa(\beta; x, y) = \frac{\exp\{-(2\kappa)^{-1} \tanh(\beta/2\kappa) (|x|^2 + |y|^2) - |x - y|^2 / (2\kappa \sinh(\beta/\kappa))\}}{\{\pi\kappa(1 - e^{-2\beta/\kappa})\}^{d/2}} . \quad (1.9)$$

Here the operator $G_\kappa(\beta)$ belongs to the *trace-class* $\mathfrak{C}_1(L^2(\mathbb{R}^d))$, with the trace-norm equals to $\text{Tr } G_\kappa(\beta) = 1/(1 - e^{-\beta/\kappa})^d = O(\kappa^d)$ for large κ . The largest eigenvalue of $G_\kappa(\beta)$ coincides with the operator norm $\|G_\kappa(\beta)\| = 1$. We write all the eigenvalues of operator $G_\kappa(\beta)$ in decreasing order:

$$g_0^{(\kappa)} = 1 > g_1^{(\kappa)} = e^{-\beta/\kappa} \geq g_2^{(\kappa)} \geq \dots .$$

The expectation value of *total* number of particles is given by

$$N_\kappa(\beta, \mu) = \frac{1}{\beta} \frac{\partial \ln \Xi_{0,\kappa}(\beta, \mu)}{\partial \mu} = \sum_{s \in \mathbb{Z}_+^d} \frac{1}{e^{\beta(\epsilon_\kappa(s) - \mu)} - 1} . \quad (1.10)$$

Since the value (1.10) diverges in the WHT limit $\kappa \rightarrow \infty$ as κ^d , one introduces the scaled quantity [DGPS], [PeSm]:

$$\rho_\kappa(\beta, \mu) := \frac{1}{\kappa^d} \sum_{s \in \mathbb{Z}_+^d} \frac{1}{e^{\beta(\epsilon_\kappa(s) - \mu)} - 1} , \quad (1.11)$$

which is a Darboux-Riemann sum for the integral

$$\rho(\beta, \mu) = \lim_{\kappa \rightarrow \infty} \rho_\kappa(\beta, \mu) = \int_{[0, \infty)^d} \frac{dp}{e^{\beta(|p|_1 - \mu)} - 1} = \sum_{s=1}^{\infty} \frac{e^{\beta \mu s}}{(\beta s)^d} . \quad (1.12)$$

Since κ^d may be interpreted as *the effective “volume”* (cf. Remark 1.1), $\rho(\beta, \mu)$ and $\rho_\kappa(\beta, \mu)$ are regarded as the expectation value of effective space-averaged density of the system, which has non-homogeneous space distribution. One defines its critical value as usual:

$$\rho_c(\beta) := \sup_{\mu < 0} \rho(\beta, \mu) = \zeta(d)/\beta^d. \quad (1.13)$$

Notice that (1.13) is bounded for $d > 1$. Therefore, if $\rho > \rho_c(\beta)$, the IBG in the WHT limit $\kappa \rightarrow \infty$ manifests a BEC in the scaled oscillator ground state (1.8) with the expected *space-averaged* condensate density:

$$\rho_0(\beta) := \rho - \rho_c(\beta) = \lim_{\kappa \rightarrow \infty} \frac{1}{\kappa^d} \frac{1}{e^{-\beta \bar{\mu}_\kappa(\beta, \rho)} - 1}. \quad (1.14)$$

Moreover the expected local density $\rho_0(\beta)(x)$ can be defined such that

$$\rho_0(\beta)(x) = \lim_{\kappa \rightarrow \infty} \frac{1}{\kappa^d} \frac{\Omega_\kappa(x)^2}{e^{-\beta \bar{\mu}_\kappa(\beta, \rho)} - 1} \quad (1.15)$$

holds. Here $\bar{\mu}_\kappa(\beta, \rho)$ is the unique root of the equation, $\rho = \rho_\kappa(\beta, \mu)$ cf. (1.11). Their limits satisfy

$$\lim_{\kappa \rightarrow \infty} \bar{\mu}_\kappa(\beta, \rho < \rho_c(\beta)) < 0 \quad \text{and} \quad \lim_{\kappa \rightarrow \infty} \bar{\mu}_\kappa(\beta, \rho \geq \rho_c(\beta)) = 0, \quad (1.16)$$

especially

$$\bar{\mu}_\kappa(\beta, \rho > \rho_c(\beta)) = -\frac{1}{\beta(\rho - \rho_c(\beta))\kappa^d} + o(\kappa^{-d}). \quad (1.17)$$

The integrated density of states $\mathcal{N}_\kappa(E)$ for the operator h_κ is given by

$$\mathcal{N}_\kappa(E) = \frac{1}{\kappa^d} \sum_{s \in \mathbb{Z}_+^d} \theta(E - |s|_1/\kappa).$$

Then from its Laplace transform

$$\int_0^\infty e^{-tE} d\mathcal{N}_\kappa(E) = [\kappa(1 - \exp(-t/\kappa))]^{-d},$$

we obtain the $\kappa \rightarrow \infty$ limit

$$d\mathcal{N}(E) = \frac{E^{d-1}}{\Gamma(d)} dE. \quad (1.18)$$

In terms of these density of states, (1.11) and (1.12) are written as

$$\rho_\kappa(\beta, \mu) = \int_0^\infty \frac{d\mathcal{N}_\kappa(E)}{e^{\beta(E-\mu)} - 1}, \quad \rho(\beta, \mu) = \int_0^\infty \frac{d\mathcal{N}(E)}{e^{\beta(E-\mu)} - 1}.$$

It is instructive to compare these results with properties of the IBG “prepared” via standard thermodynamic limit $L \rightarrow \infty$ (1.3) for e.g. Dirichlet boundary conditions. It is well-known [ZB] that in this case the expected boson density is

$$\rho_{\Lambda_L}(\beta, \mu) = \frac{1}{|\Lambda_L|} \sum_{j \in \mathbb{Z}_+} \frac{1}{e^{\beta(\varepsilon_L(j)-\mu)} - 1}. \quad (1.19)$$

Here $\{\varepsilon_L(j)\}_{j \geq 0}$ is the spectrum of the one-particle operator t_L and $\mu < \min_{j \geq 0} \varepsilon_L(j) \rightarrow 0$ as $L \rightarrow \infty$. It also has the expression

$$\rho_{\Lambda_L}(\beta, \mu) = \int_0^\infty \frac{d\tilde{\mathcal{N}}_L(E)}{e^{\beta(E-\mu)} - 1} \quad (1.20)$$

in terms of the *integrated density of states* $\{\tilde{\mathcal{N}}_L(E)\}_{L>0}$. The thermodynamic limit $\tilde{\mathcal{N}}(E) = \lim_{L \rightarrow \infty} \tilde{\mathcal{N}}_L(E)$ is independent of "non-sticky" boundary conditions [RSIV] and given by

$$d\tilde{\mathcal{N}}(E) = \frac{E^{(d-2)/2}}{(2\pi)^{d/2} \Gamma(d/2)} dE. \quad (1.21)$$

Then we get the limit of the expected density

$$\tilde{\rho}(\beta, \mu) = \lim_{L \rightarrow \infty} \rho_{\Lambda_L}(\beta, \mu) = \int_0^\infty \frac{d\tilde{\mathcal{N}}(E)}{e^{\beta(E-\mu)} - 1}. \quad (1.22)$$

The critical particle density for the IBG is

$$\tilde{\rho}_c(\beta) := \sup_{\mu < 0} \tilde{\rho}(\beta, \mu) = \int_0^\infty \frac{d\tilde{\mathcal{N}}(E)}{e^{\beta E} - 1} = \zeta(d/2)/(2\pi\beta)^{d/2}. \quad (1.23)$$

Note the difference between $\tilde{\rho}_c(\beta)$ and $\rho_c(\beta)$. In particular (1.23) is bounded only when $d > 2$. Thus thermodynamic properties of the IBG in the standard TDL $\Lambda_L \rightarrow \mathbb{R}^d$ and the WHT limit $\kappa \rightarrow \infty$ are different in spite of delusive impression that they have to produce identical systems.

Now we consider the mean-field interacting bosons trapped in the harmonic potential (1.1). Its *grand-canonical partition function* is given by

$$\Xi_{\lambda, \kappa}(\beta, \mu) := \sum_{n=0}^\infty e^{\beta(\mu n - \lambda n^2/2\kappa^d)} \text{Tr}_{\mathfrak{H}_{\text{symm}}^n} [\otimes^n G_\kappa(\beta)]. \quad (1.24)$$

We consider the case of $\beta > 0, \lambda > 0$ and arbitrary $\mu \in \mathbb{R}$. Hereafter, we suppress the symbol λ from the left-hand side of (1.24), since we fix $\lambda > 0$ in the rest of the paper.

Remark 1.1 *The scaling with the "volume" κ^d (imposed by the WHT) is a conventional way to consider the Bose-Einstein condensation in traps, see [DGPS], [LSSY], [PeSm]. Our definition of the MF interaction in WHT applies a space-average over the "volume" κ^d , which plays the same rôle as $|\Lambda_L|$ in the standard mean field model where the interaction has the form $\lambda n^2/2|\Lambda_L|$, see e.g. [ZB].*

Notice that $\lambda > 0$ corresponds to repulsive mean-field (MF) particle interaction, whereas $\lambda = 0$ is the case of the IBG (1.4).

In the present paper we consider in (1.24) only the "square" mean-field repulsive interaction $U_2 := \lambda n^2/2\kappa^d$. Although application of the Large Deviation technique makes it possible to consider also the case of the general MF interaction $U_\Phi := \kappa^d \Phi(n/\kappa^d)$, where $\Phi : \mathbb{R} \rightarrow \mathbb{R}$ is a piece-wise differentiable continuous function bounded from below, see [TZ].

To study the *non-homogeneous* condensation and the *space* distribution of the constituent bosons in the system (1.24) we use the RPF $\nu_{\kappa,\beta}$, i.e., the probability measure on the space of locally finite point measures with generating functional:

$$\begin{aligned} E_{\kappa,\beta,\mu}(f) &:= \mathbb{E}_{\kappa,\beta,\mu}[e^{-\langle f, \xi \rangle}] \\ &= \frac{1}{\Xi_{\kappa}(\beta, \mu)} \sum_{n=0}^{\infty} e^{\beta \mu n - \beta \lambda n^2 / 2\kappa^d} \text{Tr}_{\mathfrak{H}_{\text{symm}}^n} [\otimes^n (G_{\kappa}(\beta) e^{-f})], \end{aligned} \quad (1.25)$$

where $f \in C_0(\mathbb{R}^d)$, $f \geq 0$. Here $\mathbb{E}_{\kappa,\beta,\mu}[\cdot]$ stands for expectation with respect to $\nu_{\kappa,\beta,\mu}(d\xi)$, and ξ denotes the integral variable which represents locally finite point measure, see [DV, TIa, TIb, TIc]. The measure $\nu_{\kappa,\beta,\mu}$ describes a finite RPF whose *Janossy measure* can be given explicitly, see Remark 2.1.

In the present paper we study the properties of the MF interacting boson RPF $\nu_{\kappa,\beta,\mu}$ in the WHT limit $\kappa \rightarrow \infty$ by analyzing the generating functional (1.25). To this end we first define the MF *critical* chemical potential

$$\mu_{\lambda,c}(\beta) := \lambda \int_{[0,\infty)^d} \frac{dp}{e^{\beta|p|_1} - 1} = \lambda \int_0^{\infty} \frac{d\mathcal{N}(E)}{e^{\beta(E-\mu)} - 1} = \frac{\lambda \zeta(d)}{\beta^d}. \quad (1.26)$$

This critical parameter is similar to the critical chemical potential $\tilde{\mu}_c(\beta) := \lambda \tilde{\rho}_c(\beta)$ for the standard homogeneous MF boson gas via TDL, see e.g. [ZB] and (1.23).

1.3 Main Results

Now we can formulate our main theorem:

Theorem 1.2 (i) *Let $\mu < \mu_{\lambda,c}(\beta)$ (normal phase). Then the RPF $\nu_{\kappa,\beta,\mu}$ defined by (1.25), converges weakly in the WHT limit $\kappa \rightarrow \infty$ to the boson RPF ν_{β,r_*} corresponding to the generating functional:*

$$\begin{aligned} E_{\beta,r_*}(f) &:= \mathbb{E}_{\beta,r_*}[e^{-\langle f, \xi \rangle}] \\ &= \text{Det} [1 + \sqrt{1 - e^{-f}} r_* G(\beta) (1 - r_* G(\beta))^{-1} \sqrt{1 - e^{-f}}]^{-1}. \end{aligned} \quad (1.27)$$

Here $\mathbb{E}_{\beta,r_*}[\cdot]$ denotes expectation with respect to the measure ν_{β,r_*} , Det stands for the Fredholm determinant, $G(\beta) := e^{\beta \Delta/2}$ is the heat semigroup on \mathfrak{H} and $r_* = r_*(\beta, \mu, \lambda) \in (0, 1)$ is a unique solution of the equation :

$$\beta \mu = \log r + \lambda \beta \int_0^{\infty} \frac{d\mathcal{N}(E)}{r^{-1} e^{\beta E} - 1}. \quad (1.28)$$

(ii) *For $\mu > \mu_{\lambda,c}(\beta)$ (condensed phase), the generating functional (1.25) has the following asymptotics:*

$$\lim_{\kappa \rightarrow \infty} \frac{1}{\kappa^{d/2}} \log E_{\kappa,\beta,\mu}(f) = - \frac{\mu - \mu_{\lambda,c}(\beta)}{\pi^{d/2} \lambda} (\sqrt{1 - e^{-f}}, (1 + K_f)^{-1} \sqrt{1 - e^{-f}}), \quad (1.29)$$

where $K_f := \left(G(\beta)^{1/2} (1 - G(\beta))^{-1/2} \sqrt{1 - e^{-f}} \right)^* \left(G(\beta)^{1/2} (1 - G(\beta))^{-1/2} \sqrt{1 - e^{-f}} \right)$ is a positive trace-class operator on $\mathfrak{H} = L^2(\mathbb{R}^d)$ for $d > 2$.

Remark 1.3 For dimensions $d > 1$ the integral (1.26) is finite. This gives an idea that the BEC is possible for the case $\mu > \mu_{\lambda,c}(\beta)$ and dimensions $d > 1$. However, in the present paper we assume $d > 2$ to be able to prove our main theorem. In fact for $d = 2$, the operator K_f is not a trace-class operator and the Fredholm determinant is not well-defined, see [Tib].

Remark 1.4 Because of a technical difficulty, actually we do not have results for the critical case $\mu = \mu_{\lambda,c}(\beta)$.

Remark 1.5 The heat semigroup $G(\beta)$ is appeared here since its kernel

$$G(\beta; x, y) = (2\pi\beta)^{-d/2} e^{-|x-y|^2/2\beta}$$

is the point-wise limit of the Mehler kernel (1.9), as $\kappa \rightarrow \infty$. Therefore the generating functional of the resulting RPF (1.27) in (i) has exactly the same form as that for the standard homogeneous IBG in the non-condensed phase, see eq.(2.13) in [Tia], see also [ShTa]. That is, the position distribution of the model in the non-condensed phase coincides with that of the standard IBG. However, it is the integrated density of states $\mathcal{N}(E)$ that is appeared in (1.28), instead of $\tilde{\mathcal{N}}(E)$. It implies that the dependence of r_* as a function of parameters, especially β , is different from standard IBG or the standard MF-model.

Remark 1.6 The RPF describing BEC for homogeneous IBG is given by the convolution of two RPFs [Tib]. One convolution component is the usual boson RPF, while the other component seems to describe the position distribution of the condensed part of the constituent bosons. The behaviour of the generating functional in case (ii) shows that the latter component overwhelms the former in the present model. It is to be noted here that the latter can not be explained by the particles in the ground state alone, it contains the effect of the interference between “the condensed part” and “the normal part” [EK], although the intensity of the RPF is proportional to the square of the ground state wave function, as (1.15).

The sharp contrast between two regimes (i) and (ii) in Theorem 1.2 may be seen by the expectation values.

Corollary 1.7 For the case (i) $\mu < \mu_{\lambda,c}(\beta)$ (normal phase)

$$\mathbb{E}_{\beta,r_*}[\langle f, \xi \rangle] = \text{Tr} [f r_* G(\beta)(1 - r_* G(\beta))^{-1}] = \rho_{r_*} \int_{\mathbb{R}^d} dx f(x)$$

holds, where ρ_{r_*} is given by

$$\rho_{r_*} = r_* G(\beta)(1 - r_* G(\beta))^{-1}(x, x) = \sum_{n=1}^{\infty} r_*^n / (2\pi\beta n)^{d/2}.$$

For the case (ii) $\mu > \mu_{\lambda,c}(\beta)$ (condensed phase),

$$\liminf_{\kappa \rightarrow \infty} \frac{\mathbb{E}_{\kappa,\beta,\mu,\lambda}[\langle f, \xi \rangle]}{\kappa^{d/2}} \geq \frac{\mu - \mu_{\lambda,c}(\beta)}{\pi^{d/2}\lambda} \int_{\mathbb{R}^d} dx f(x) \quad (1.30)$$

holds, where $f \in C_0(\mathbb{R}^d)$.

The *weak limits* of the RPFs concerns the limit of the *local* position distribution of particles. In this sense, the results of Theorem 1.2 and Corollary 1.7 in regime (i) may be interpreted as follows: in the WHT limit the position distribution of the MF interacting bosons in neighbourhoods of the origin of coordinates (i.e. the bottom of the WHT potential) is close to that of a free IBG corresponding to the unconventional parameter (1.28). The information about the particle position distribution in domains distant from the bottom of the WHT are missing in the limit ν_{β, r_*} . In order to take this “tail” particles into account we use the standard definition of the grand-canonical total number of particles for our model :

$$\begin{aligned} \rho_{\kappa, \lambda}^{(tot)}(\beta, \mu) &:= \frac{1}{\kappa^d \beta} \frac{\partial \ln \Xi_{\kappa}(\beta, \mu)}{\partial \mu} \\ &= \frac{1}{\kappa^d \Xi_{\kappa, \lambda}(\beta, \mu)} \sum_{n=0}^{\infty} n e^{\beta(\mu n - \lambda n^2 / 2\kappa^d)} \text{Tr}_{\mathfrak{H}_{sym}^n} [\otimes^n G_{\kappa}(\beta)] . \end{aligned} \quad (1.31)$$

Since κ^d is interpreted as the effective volume of the model, $\rho_{\kappa, \lambda}^{(tot)}(\beta, \mu)$ represents the effective total *space-averaged* density of the non-homogeneous system (1.24).

Theorem 1.8 *The WHT limit of (1.31)*

$$\rho_{\lambda}^{(tot)}(\beta, \mu) = \lim_{\kappa \rightarrow \infty} \rho_{\kappa, \lambda}^{(tot)}(\beta, \mu) = \lim_{\kappa \rightarrow \infty} \kappa^{-d} \text{Tr} [r_* G_{\kappa} (1 - r_* G_{\kappa})^{-1}] \quad (1.32)$$

exists and satisfies :

(i) ($\mu \leq \mu_{\lambda, c}(\beta)$)

$$\rho_{\lambda}^{(tot)}(\beta, \mu) = \int_{[0, \infty)^d} \frac{d\mathcal{N}(E)}{r_*^{-1} e^{\beta E} - 1} \quad \text{and} \quad \beta \mu = \log r_* + \lambda \beta \rho_{\lambda}^{(tot)}(\beta, \mu) ; \quad (1.33)$$

(ii) ($\mu > \mu_{\lambda, c}(\beta)$)

$$\rho_{\lambda}^{(tot)}(\beta, \mu) = \mu / \lambda . \quad (1.34)$$

It also holds that

$$\rho_c^{(tot)}(\beta) := \lim_{\mu \rightarrow \mu_c(\beta)} \rho_{\lambda}^{(tot)}(\beta, \mu) = \zeta(d) / \beta^d . \quad (1.35)$$

Remark 1.9 *The readers should not to confuse two “densities” : ρ_{r_*} in Corollary 1.7 and $\rho_{\lambda}^{(tot)}(\beta, \mu)$ defined above. The ρ_{r_*} can be interpreted as the limit of the “local” density around the origin of coordinate of non-homogeneous RPF $\nu_{\kappa, \beta, \mu}$, on the other hand the $\rho_{\lambda}^{(tot)}(\beta, \mu)$ retains the information about the expectation of the total number of particles with respect to $\nu_{\kappa, \beta, \mu}$ through the WHT limit.*

Note that $\rho_c^{(tot)}(\beta)$ coincides with (1.13) of IBG in the WHT.

Remark 1.10 *Qualitatively different behaviour of the space distributions of bosons described in Theorem 1.2 can be understand heuristically with the help of Theorem 1.8 in the following way. Consider the WHT limit $\kappa \rightarrow \infty$:*

In case (i) the bosons are distributed almost uniformly in the region of radius κ according to the kernel (1.9).

On the other hand, in case (ii) (condensed phase) the condensed part of particles $\kappa^d(\rho_\lambda^{(tot)}(\beta, \mu) - \rho_{\lambda,c}^{(tot)}(\beta)) = \kappa^d(\mu - \mu_{\lambda,c}(\beta))/\lambda$ is localized in the region of radius $O(\kappa^{1/2})$ according to the profile of the square of the ground state wave function Ω_κ . Whereas the particles outside of the condensate essentially spread out over the region of radius κ .

The paper is organized as follows. Preliminary estimates and results concerning the WHT limit for the *mean-field* interacting boson gas ($\lambda > 0$) are collected in Section 2. Section 3 and 4 are dedicated to the proof of Theorem 1.2 and 1.8, respectively. We reserved Section 5 for summary and conjectures.

2 Preliminary Arguments and Estimates

In this section, we write the expectation (1.25) as the ratio $\tilde{\Xi}_\kappa(\beta, \mu)/\Xi_\kappa(\beta, \mu)$. The representations of $\tilde{\Xi}_\kappa(\beta, \mu)$ and $\Xi_\kappa(\beta, \mu)$ are given in the form of integration of Fredholm determinants. We also give the miscellaneous estimates needed for the evaluation of these integrals.

2.1 $\Xi_\kappa(\beta, \mu)$ and $\tilde{\Xi}_\kappa(\beta, \mu)$

In terms of the projection operator on $\mathfrak{H}^n = \otimes^n L^2(\mathbb{R}^d)$ onto its subspace \mathfrak{H}_{sym}^n , the grand-canonical partition function can be written as

$$\Xi_\kappa(\beta, \mu) = \sum_{n=0}^{\infty} e^{\beta\mu n - \beta\lambda n^2/2\kappa^d} \frac{1}{n!} \sum_{\sigma \in \mathcal{S}_n} \text{Tr}_{\otimes^n L^2(\mathbb{R}^d)} [(\otimes^n G_\kappa(\beta))U(\sigma)],$$

where the second sum is taken over the symmetric group \mathcal{S}_n and

$$U(\sigma)\varphi_1 \otimes \cdots \otimes \varphi_n = \varphi_{\sigma^{-1}(1)} \otimes \cdots \otimes \varphi_{\sigma^{-1}(n)} \quad \text{for } \sigma \in \mathcal{S}_n, \varphi_1, \dots, \varphi_n \in L^2(\mathbb{R}^d).$$

Hence we have

$$\begin{aligned} \Xi_\kappa(\beta, \mu) &= \sum_{n=0}^{\infty} \frac{e^{\beta\mu n - \beta\lambda n^2/2\kappa^d}}{n!} \sum_{\sigma \in \mathcal{S}_n} \int_{(\mathbb{R}^d)^n} \left(\prod_{j=1}^n G_\kappa(\beta)(x_j, x_{\sigma^{-1}(j)}) \right) dx_1 \cdots dx_n \\ &= \sum_{n=0}^{\infty} \frac{e^{\beta\mu n - \beta\lambda n^2/2\kappa^d}}{n!} \int_{(\mathbb{R}^d)^n} \text{Per} \{G_\kappa(\beta)(x_i, x_j)\}_{1 \leq i, j \leq n} dx_1 \cdots dx_n, \end{aligned}$$

here Per stands for the permanent of the matrix $\{G_\kappa(\beta)(x_i, x_j)\}_{1 \leq i, j \leq n}$.

Remark 2.1 The point field $\nu_{\kappa, \beta, \mu}$ of (1.25) can also be defined in terms of Janossy measures or exclusion probability [DV]. This means that $\nu_{\kappa, \beta, \mu}$ is a finite point field, which assigns the probability

$$\text{Pr}\{dX_n\} := \frac{e^{\beta\mu n - \beta\lambda n^2/2\kappa^d}}{n! \Xi_\kappa(\beta, \mu)} \text{Per} \{G_\kappa(\beta)(x_i, x_j)\}_{1 \leq i, j \leq n} dx_1 \cdots dx_n$$

to the event $\{dX_n\}$: there are exactly n points, one in each infinitesimal region $\prod_{i=1}^d (x_j^{(i)}, x_j^{(i)} + dx_j^{(i)})$, $(x_j = (x_j^{(1)}, \dots, x_j^{(d)}), j = 1, \dots, n)$.

As in [TIa, TIb], we use the generalized *Vere-Jones' formula* [ShTa, V] in the form

$$\frac{1}{n!} \int Per \{J(x_i, x_j)\}_{1 \leq i, j \leq n} dx_1 \cdots dx_n = \oint_{S_r(0)} \frac{dz}{2\pi i z^{n+1} \text{Det}(1 - zJ)},$$

where $r > 0$ satisfies $\|rJ\| < 1$. $S_r(\zeta)$ denotes the integration contour defined by the map $\theta \mapsto \zeta + r \exp(i\theta)$, where θ ranges from $-\pi$ to π , $r > 0$ and $\zeta \in \mathbb{C}$. Then we obtain

$$\Xi_\kappa(\beta, \mu) = \sum_{n=0}^{\infty} e^{\beta\mu n - \beta\lambda n^2/2\kappa^d} \oint_{S_r(0)} \frac{dz}{2\pi i z^{n+1} \text{Det}(1 - zG_\kappa(\beta))},$$

where $r \in (0, \|G_\kappa(\beta)\|^{-1}) = (0, 1)$. Note that the zeroth term is 1 in this expression.

Let us substitute

$$e^{-\beta\lambda n^2/2\kappa^d} = \sqrt{\frac{\beta\lambda}{2\pi\kappa^d}} \int_{\mathbb{R}} dx \exp\left(-\frac{\beta\lambda}{2\kappa^d}((x + is)^2 - 2in(x + is))\right). \quad (2.1)$$

If $s > 0$ satisfies

$$e^{\beta\mu - \beta\lambda s/\kappa^d} < r,$$

we can take the summation over n together with the complex integration and a scaling of x to get

$$\begin{aligned} \Xi_\kappa(\beta, \mu) &= \sqrt{\frac{\beta\lambda}{2\pi\kappa^d}} e^{\beta\lambda s^2/2\kappa^d} \int_{\mathbb{R}} dx \oint \frac{dz}{2\pi i} \frac{e^{-\beta\lambda(x^2 + 2isx)/2\kappa^d}}{(z - e^{\beta\mu + \beta\lambda(ix - s)/\kappa^d}) \text{Det}[1 - zG_\kappa(\beta)]} \\ &= \sqrt{\frac{\kappa^d}{2\pi\beta\lambda}} e^{\beta\lambda s^2/2\kappa^d} \int_{\mathbb{R}} dx \frac{e^{-isx - \kappa^d x^2/2\beta\lambda}}{\text{Det}[1 - e^{\beta\mu + ix - \beta\lambda s/\kappa^d} G_\kappa(\beta)]}. \end{aligned} \quad (2.2)$$

Note that after z -integration, r disappears and (2.2) is valid for any s satisfying $\exp(\beta\mu - \beta\lambda s/\kappa^d) \in (0, \|G_\kappa(\beta)\|^{-1})$. We will estimate the integral in the spirit of saddle point method. Here, we extract the main part from the integral. Let $s = s_\kappa, r = r_\kappa$ be the unique solution of the system:

$$\begin{cases} r = \exp(\beta\mu - \beta\lambda s/\kappa^d) \\ s = \text{Tr}[rG_\kappa(\beta)(1 - rG_\kappa(\beta))^{-1}]. \end{cases} \quad (2.3)$$

Obviously, the condition $r_\kappa \in (0, \|G_\kappa(\beta)\|^{-1})$ is fulfilled. Hence, we can substitute in (2.2) s by s_κ . Using the product property of the Fredholm determinant, we get for denominator of (1.25) the representation:

$$\Xi_\kappa(\beta, \mu) = \sqrt{\frac{\kappa^d}{2\pi\beta\lambda}} \frac{e^{\beta\lambda s_\kappa^2/2\kappa^d}}{\text{Det}[1 - r_\kappa G_\kappa(\beta)]} \int_{\mathbb{R}} dx \frac{e^{-is_\kappa x - \kappa^d x^2/2\beta\lambda}}{\text{Det}[1 - (e^{ix} - 1)r_\kappa G_\kappa(\beta)(1 - r_\kappa G_\kappa(\beta))^{-1}]} \quad (2.4)$$

For the numerator of (1.25), we introduce bounded symmetric operators

$$\tilde{G}_\kappa(\beta)(f) := G_\kappa(\beta)^{1/2} e^{-f} G_\kappa(\beta)^{1/2}, \quad (2.5)$$

indexed by function $f \in C_0(\mathbb{R}^d)$, $f \geq 0$, which we skip below for simplicity. Then for generating functional (1.25) one gets the form: $E_{\kappa,\beta,\mu}(f) = \tilde{\Xi}_\kappa(\beta, \mu)(f) / \Xi_\kappa(\beta, \mu)$, where

$$\begin{aligned} \tilde{\Xi}_\kappa(\beta, \mu)(f) &= \tilde{\Xi}_\kappa(\beta, \mu) = \sum_{n=0}^{\infty} e^{\beta\mu n - \beta\lambda n^2 / 2\kappa^d} \text{Tr}_{\mathfrak{H}_{sym}^n} [\otimes^n \tilde{G}_\kappa(\beta)] \\ &= \sqrt{\frac{\kappa^d}{2\pi\beta\lambda}} \frac{e^{\beta\lambda\tilde{s}_\kappa^2/2\kappa^d}}{\text{Det}[1 - \tilde{r}_\kappa \tilde{G}_\kappa(\beta)]} \int_{\mathbb{R}} dx \frac{e^{-i\tilde{s}_\kappa x - \kappa^d x^2 / 2\beta\lambda}}{\text{Det}[1 - (e^{ix} - 1)\tilde{r}_\kappa \tilde{G}_\kappa(\beta)(1 - \tilde{r}_\kappa \tilde{G}_\kappa(\beta))^{-1}]}. \end{aligned} \quad (2.6)$$

Here $(\tilde{s}_\kappa, \tilde{r}_\kappa)$ is the unique solution of

$$\begin{cases} \tilde{r} = \exp(\beta\mu - \beta\lambda\tilde{s}/\kappa^d) \\ \tilde{s} = \text{Tr}[\tilde{r}\tilde{G}_\kappa(\beta)(1 - \tilde{r}\tilde{G}_\kappa(\beta))^{-1}]. \end{cases} \quad (2.7)$$

Obviously, $\tilde{r}_\kappa \in (0, \|\tilde{G}_\kappa(\beta)\|^{-1})$. Note also that r_κ and \tilde{r}_κ satisfy the following conditions respectively:

$$\frac{1}{\kappa^d} \text{Tr}[r_\kappa G_\kappa(\beta)(1 - r_\kappa G_\kappa(\beta))^{-1}] = \frac{\beta\mu - \log r_\kappa}{\beta\lambda}, \quad (2.8)$$

$$\frac{1}{\kappa^d} \text{Tr}[\tilde{r}_\kappa \tilde{G}_\kappa(\beta)(1 - \tilde{r}_\kappa \tilde{G}_\kappa(\beta))^{-1}] = \frac{\beta\mu - \log \tilde{r}_\kappa}{\beta\lambda}. \quad (2.9)$$

Since by definition (2.5) one obviously gets: $\tilde{G}_\kappa(\beta) \leq G_\kappa(\beta)$, the operator $\tilde{G}_\kappa(\beta)$ also belongs to the trace-class $\mathcal{C}_1(\mathfrak{H})$. We put the eigenvalues of $\tilde{G}_\kappa(\beta)$ in the decreasing order

$$\tilde{g}_0^{(\kappa)} = \|\tilde{G}_\kappa(\beta)\| \geq \tilde{g}_1^{(\kappa)} \geq \dots$$

Then, we have $g_j^{(\kappa)} \geq \tilde{g}_j^{(\kappa)}$ ($j = 0, 1, 2, \dots$) by the *min-max principle*.

2.2 Approximations of One-Particle Gibbs Semigroups

Here we establish some relations between Gibbs semigroup $\{G_\kappa(\beta)\}_{\beta \geq 0}$ and the heat semigroup $\{G(\beta)\}_{\beta \geq 0}$. Let P_κ be the orthogonal projection on \mathfrak{H} onto its one-dimensional subspace spanned by the vector Ω_κ , and put $Q_\kappa := I - P_\kappa$.

Lemma 2.2 *For any $r \in (0, 1)$,*

$$\|\sqrt{1 - e^{-f}} [rG_\kappa(\beta)(1 - rG_\kappa(\beta))^{-1} - rG(\beta)(1 - rG(\beta))^{-1}] \sqrt{1 - e^{-f}}\|_1 \rightarrow 0, \quad (2.10)$$

$$\|\sqrt{1 - e^{-f}} Q_\kappa G_\kappa(\beta) Q_\kappa (1 - Q_\kappa G_\kappa(\beta) Q_\kappa)^{-1} \sqrt{1 - e^{-f}} - K_f\|_1 \rightarrow 0 \quad (2.11)$$

hold in the limit $\kappa \rightarrow \infty$, where $\|\cdot\|_1$ stands for the trace norm in $\mathcal{C}_1(L^2(\mathbb{R}^d))$.

Proof : First, we show the estimates

$$|G_\kappa^n(\beta; x, y) - G^n(\beta; x, y)| \leq \frac{A'}{\kappa n^{d/2-1}} \left(1 + \frac{|x|^2 + |y|^2}{\kappa}\right) \quad \text{if } n\beta/\kappa \leq 1, \quad (2.12)$$

$$|G_\kappa^n(\beta; x, y) - \Omega_\kappa(x)\Omega_\kappa(y)| \leq \frac{B'e^{-n\beta/2\kappa}}{\kappa^{d/2}} \left(1 + \frac{|x-y|^2}{\kappa}\right) \quad \text{if } n\beta/\kappa \geq 1, \quad (2.13)$$

where A' and B' depend only on d and β . In fact, by the *Mehler's formula* one gets for $n\beta/\kappa \leq 1$ the estimate:

$$\begin{aligned} & |G_\kappa^n(\beta; x, y) - G^n(\beta; x, y)| \\ & \leq \frac{1}{(2\pi n\beta)^{d/2}} \left| \left(\frac{2n\beta/\kappa}{1 - e^{-2n\beta/\kappa}} \right)^{d/2} - 1 \right| \exp \left(- \frac{\tanh(n\beta/2\kappa)}{2\kappa} (|x|^2 + |y|^2) - \frac{|x-y|^2}{2\kappa \sinh(n\beta/\kappa)} \right) \\ & + \frac{1}{(2\pi n\beta)^{d/2}} \left| \exp \left(- \frac{\tanh(n\beta/2\kappa)}{2\kappa} (|x|^2 + |y|^2) \right) - 1 \right| \exp \left(- \frac{|x-y|^2}{2\kappa \sinh(n\beta/\kappa)} \right) \\ & + \frac{1}{(2\pi n\beta)^{d/2}} \left| 1 - \exp \left(- \left(\frac{\sinh(n\beta/\kappa)}{n\beta/\kappa} - 1 \right) \frac{|x-y|^2}{2\kappa \sinh(n\beta/\kappa)} \right) \right| \exp \left(- \frac{|x-y|^2}{2\kappa \sinh(n\beta/\kappa)} \right) \\ & \leq \frac{1}{(2\pi n\beta)^{d/2}} \left(\frac{An\beta}{\kappa} + \frac{|x|^2 + |y|^2}{2\kappa} \frac{n\beta}{2\kappa} + \left(\frac{n\beta}{\kappa} \right)^2 \frac{|x-y|^2}{2\kappa \sinh(n\beta/\kappa)} \right) \exp \left(- \frac{|x-y|^2}{2\kappa \sinh(n\beta/\kappa)} \right) \\ & \leq \frac{A'}{\kappa n^{d/2-1}} \left(1 + \frac{|x|^2 + |y|^2}{\kappa} \right). \end{aligned}$$

Here we have used (A.2) for the first term, (A.4, A.6) for the second term and (A.4, A.8) for the third term at the second inequality and (A.8) at the last inequality.

On the other hand for $n\beta/\kappa \geq 1$, we obtain:

$$\begin{aligned} & |G_\kappa^n(\beta; x, y) - \Omega_\kappa(x)\Omega_\kappa(y)| \\ & \leq \frac{1}{(\pi\kappa)^{d/2}} \left| \left(\frac{1}{1 - e^{-2n\beta/\kappa}} \right)^{d/2} - 1 \right| \exp \left(- \frac{\tanh(n\beta/2\kappa)}{2\kappa} (|x|^2 + |y|^2) - \frac{|x-y|^2}{2\kappa \sinh(n\beta/\kappa)} \right) \\ & + \frac{1}{(\pi\kappa)^{d/2}} \left| \exp \left(- \frac{\tanh(n\beta/2\kappa)}{2\kappa} (|x|^2 + |y|^2) \right) - \exp \left(- \frac{|x|^2 + |y|^2}{2\kappa} \right) \right| \exp \left(- \frac{|x-y|^2}{2\kappa \sinh(n\beta/\kappa)} \right) \\ & + \frac{1}{(\pi\kappa)^{d/2}} \exp \left(- \frac{|x|^2 + |y|^2}{2\kappa} \right) \left| \exp \left(- \frac{|x-y|^2}{2\kappa \sinh(n\beta/\kappa)} \right) - 1 \right| \\ & \leq \frac{1}{(\pi\kappa)^{d/2}} \left(B e^{-2n\beta/\kappa} + \frac{1}{e} \left(\coth \left(\frac{n\beta}{2\kappa} \right) - 1 \right) + \frac{|x-y|^2}{2\kappa \sinh(n\beta/\kappa)} \right) \\ & \leq \frac{B'e^{-n\beta/2\kappa}}{\kappa^{d/2}} \left(1 + \frac{|x-y|^2}{\kappa} \right), \end{aligned}$$

where we have used (A.3) for the first term, (A.5, A.7) for the second term and (A.4, A.9) for the third term.

Now, let us show the second limit (2.11), notice that the inequality:

$$\begin{aligned}
& \sum_{n=1}^{\infty} |G_{\kappa}^n(\beta; x, y) - G^n(\beta; x, y) - \Omega_0^{\kappa}(x)\Omega_{0,\kappa}(y)| \\
& \leq \sum_{n=1}^{\lceil \kappa/\beta \rceil} \left(\frac{A'}{\kappa n^{d/2-1}} \left(1 + \frac{|x|^2 + |y|^2}{\kappa} \right) + \frac{1}{(\pi\kappa)^{d/2}} \right) \\
& + \sum_{n=\lceil \kappa/\beta \rceil + 1}^{\infty} \left(\frac{B' e^{-n\beta/2\kappa}}{\kappa^{d/2}} \left(1 + \frac{|x-y|^2}{\kappa} \right) + \frac{1}{(2\pi n\beta)^{d/2}} \right) \\
& \leq C'(\kappa^{1-d/2} \vee \kappa^{-1} \log \kappa) \left(1 + \frac{|x|^2 + |y|^2}{\kappa} \right),
\end{aligned}$$

holds for C' , which depends only on d and β . The *integer part* is denoted by $\lceil \cdot \rceil$. Here we used the estimates (2.12), (2.13) and

$$\sum_{n=1}^N \frac{1}{n^{d/2-1}} = \begin{cases} O(1) & (d/2 > 2) \\ O(N^{2-d/2}) & (1 < d/2 < 2) \\ O(\log N) & d/2 = 2 \end{cases}$$

and so on. Now put

$$A^{(\kappa)} := \sqrt{1 - e^{-f}} Q_{\kappa} G_{\kappa}(\beta) Q_{\kappa} (1 - Q_{\kappa} G_{\kappa}(\beta) Q_{\kappa})^{-1} \sqrt{1 - e^{-f}}.$$

Then, since $\|Q_{\kappa} G_{\kappa}(\beta) Q_{\kappa}\| = e^{-\beta/\kappa} < 1$, one gets in the limit $N \rightarrow \infty$ the operator-norm convergence:

$$A_N^{(\kappa)} = \sqrt{1 - e^{-f}} \sum_{n=1}^N Q_{\kappa} G_{\kappa}(\beta)^n Q_{\kappa} \sqrt{1 - e^{-f}} \rightarrow A^{(\kappa)} \geq 0. \quad (2.14)$$

Recall that Theorem 3.1(i) and Proposition 2.3(i) of [Tib] yield the *strong* convergence:

$$K_N = \sqrt{1 - e^{-f}} \sum_{n=1}^N G(\beta)^n \sqrt{1 - e^{-f}} \rightarrow K_f \geq 0 \quad (2.15)$$

for $N \rightarrow \infty$. Moreover, we also have the following estimate for the operator norm.

$$\begin{aligned}
\|A_N^{(\kappa)} - K_N\| &= \sup_{\|\phi\|=1} |(\phi, \sqrt{1 - e^{-f}} \sum_{n=1}^N (Q_{\kappa} G_{\kappa}(\beta)^n Q_{\kappa} - G(\beta)^n) \sqrt{1 - e^{-f}} \phi)| \\
&\leq \sup_{\|\phi\|=1} \int_{\text{supp } f} dx \int_{\text{supp } f} dy \sqrt{1 - e^{-f(x)}} \sqrt{1 - e^{-f(y)}} |\phi(x)\phi(y)| \\
&\quad \times \sum_{n=1}^N |G_{\kappa}^n(\beta; x, y) - G^n(\beta; x, y) - \Omega_{\kappa}(x)\Omega_{\kappa}(y)|
\end{aligned}$$

$$\leq \|\sqrt{1-e^{-f}}\|^2 \sup_{x,y \in \text{supp } f} \sum_{n=1}^{\infty} |G_{\kappa}^n(\beta; x, y) - G^n(\beta; x, y) - \Omega_0^{\kappa}(x)\Omega_{0,\kappa}(y)| \rightarrow 0$$

for $\kappa \rightarrow \infty$ *uniformly* in N . Here $\|\sqrt{1-e^{-f}}\|$ stands for the L^2 norm of the function. We have used Cauchy-Schwarz inequality at the second inequality. The standard 3ϵ -argument yields that $A^{(\kappa)} \rightarrow K_f$ strongly when $\kappa \rightarrow \infty$.

On the other hand, since the operators $A^{(\kappa)}, K_f$ are *non-negative*, we have for $\kappa \rightarrow \infty$ the limit:

$$\begin{aligned} \|A^{(\kappa)}\|_1 - \|K_f\|_1 &= \text{Tr } A^{(\kappa)} - \text{Tr } K_f = \sum_{n=1}^{\infty} (\phi_n, (A^{(\kappa)} - K_f) \phi_n) \\ &= \sum_{n=1}^{\infty} \sum_{l=1}^{\infty} (\phi_n, \sqrt{1-e^{-f}} (Q_{\kappa} G_{\kappa}(\beta)^l Q_{\kappa} - G(\beta)^l) \sqrt{1-e^{-f}} \phi_n) \\ &= \sum_{l=1}^{\infty} \sum_{n=1}^{\infty} (\phi_n, \sqrt{1-e^{-f}} (Q_{\kappa} G_{\kappa}(\beta)^l Q_{\kappa} - G(\beta)^l) \sqrt{1-e^{-f}} \phi_n) \\ &= \sum_{l=1}^{\infty} \text{Tr } \sqrt{1-e^{-f}} (Q_{\kappa} G_{\kappa}(\beta)^l Q_{\kappa} - G(\beta)^l) \sqrt{1-e^{-f}} \\ &= \sum_{l=1}^{\infty} \int_{\text{supp } f} (1 - e^{-f(x)}) (G_{\kappa}^l(\beta; x, x) - G^l(\beta; x, x) - \Omega_{\kappa}(x)^2) dx \rightarrow 0, \end{aligned}$$

where $\{\phi_n\}_{n=1}^{\infty}$ is an arbitrary complete ortho-normal system in \mathfrak{H} . Note that we can exchange the order of summations over n and l , since

$$\text{Tr } A^{(\kappa)} = \sum_{n,l} (\phi_n, \sqrt{1-e^{-f}} Q_{\kappa} G_{\kappa}(\beta)^l Q_{\kappa} \sqrt{1-e^{-f}} \phi_n)$$

and

$$\text{Tr } K_f = \sum_{n,l} (\phi_n, \sqrt{1-e^{-f}} G^l(\beta) \sqrt{1-e^{-f}} \phi_n)$$

are convergent non-negative sequences. Here we used (2.14) and (2.15) in the third equality above. Thus, we get $\lim_{\kappa \rightarrow \infty} A^{(\kappa)} = K_f$ in $\mathfrak{C}_1(\mathfrak{H})$ by the *Grümm convergence* theorem, see e.g. [Z].

Let us consider the first limit (2.10). Since the identity $rG_{\kappa}(\beta)(1-rG_{\kappa}(\beta))^{-1} = \sum_{n=1}^{\infty} r^n G_{\kappa}(\beta)^n$ holds in the operator norm topology and $G_{\kappa}^n(\beta; x, y) \geq 0$, (by virtue of Lemma 2.2(ii) in [Tib]) we get the representation:

$$[rG_{\kappa}(\beta)(1-rG_{\kappa}(\beta))^{-1}](x, y) = \sum_{n=1}^{\infty} r^n G_{\kappa}^n(\beta; x, y).$$

Similarly, one gets the representation:

$$[rG(\beta)(1-rG(\beta))^{-1}](x, y) = \sum_{n=1}^{\infty} r^n G^n(\beta; x, y).$$

In fact the series in the right-hand side of the above representations are *point-wise* convergent because the uniform estimates

$$G_\kappa^n(\beta; x, y) \leq \left(\frac{1}{\pi\beta} \vee \frac{2}{\pi\kappa} \right)^{d/2}, \quad G^n(\beta; x, y) \leq \left(\frac{1}{2\pi\beta} \right)^{d/2}$$

hold for all $\kappa > 0$, $x, y \in \mathbb{R}^d$ and $n \in \mathbb{N}$. Here we used (A.1) for the first inequality. Hence we obtain the estimate

$$\begin{aligned} & \left| [rG_\kappa(\beta)(1-rG_\kappa(\beta))^{-1}](x, y) - [rG(\beta)(1-rG(\beta))^{-1}](x, y) \right| \leq \sum_{n=1}^{\infty} r^n |G_\kappa^n(\beta; x, y) - G^n(\beta; x, y)| \\ & \leq \sum_{n=1}^{\lceil \kappa/\beta \rceil} r^n \frac{A'}{\kappa n^{d/2-1}} \left(1 + \frac{|x|^2 + |y|^2}{\kappa} \right) + \sum_{n=\lceil \kappa/\beta \rceil+1}^{\infty} r^n \left[\left(\frac{1}{2\pi\beta} \right)^{d/2} + \left(\frac{1}{\pi\beta} \vee \frac{2}{\pi\kappa} \right)^{d/2} \right] \\ & \leq \frac{A'}{\kappa(1-r)} \left(1 + \frac{|x|^2 + |y|^2}{\kappa} \right) + C \frac{r^{\lceil \kappa/\beta \rceil+1}}{1-r} \end{aligned}$$

which tends to zero when $\kappa \rightarrow \infty$, uniformly in $x, y \in \mathfrak{C}$ for any compact set \mathfrak{C} , where C denotes a constant which depends only on d and on β .

Thus,

$$\sup_{x, y \in \text{supp } f} \left| [rG_\kappa(\beta)(1-rG_\kappa(\beta))^{-1}](x, y) - [rG(\beta)(1-rG(\beta))^{-1}](x, y) \right| \rightarrow 0$$

holds. We get the first of the announced limits (2.10) by the similar (even simpler) argument to the second one. This finishes the proof of the lemma. \square

2.3 Estimates for the Scaled Mean-Field Interaction

In the followings, we use the notation $B_\kappa := \hat{O}(\kappa^\alpha)$, which means that there exist two numbers $c_1 \geq c_2 > 0$ such that

$$c_1 \kappa^\alpha \geq B_\kappa \geq c_2 \kappa^\alpha.$$

Next, we put $W_\kappa := (G_\kappa(\beta))^{1/2} \sqrt{1-e^{-f}}$ and define $D_\kappa := G_\kappa(\beta) - \tilde{G}_\kappa(\beta) = W_\kappa W_\kappa^*$.

Lemma 2.3 *For large $\kappa > 0$, the following asymptotics hold:*

$$\begin{aligned} (\Omega_\kappa, D_\kappa \Omega_\kappa) &= \|W_\kappa^* \Omega_\kappa\|^2 = \|\sqrt{1-e^{-f}}\|^2 (1+o(1)) / (\pi\kappa)^{d/2}, \\ \text{Tr } D_\kappa &= \frac{\|\sqrt{1-e^{-f}}\|^2 (1+o(1))}{(\pi\kappa(1-e^{-2\beta/\kappa}))^{d/2}} \end{aligned}$$

and

$$\begin{aligned} g_0^{(\kappa)} - \tilde{g}_0^{(\kappa)} &= (\Omega_\kappa, (D_\kappa - D_\kappa Q_\kappa [\tilde{g}_0^{(\kappa)} - Q_\kappa \tilde{G}_\kappa(\beta) Q_\kappa]^{-1} Q_\kappa D_\kappa) \Omega_\kappa) \\ &= \frac{1+o(1)}{(\pi\kappa)^{d/2}} (\sqrt{1-e^{-f}}, [1 + W_\kappa^* Q_\kappa [1 - Q_\kappa G_\kappa(\beta) Q_\kappa]^{-1} Q_\kappa W_\kappa]^{-1} \sqrt{1-e^{-f}}) \\ &= \hat{O}(\kappa^{-d/2}). \end{aligned}$$

Proof: For simplicity of notation we suppress everywhere below the index κ in $g_j^{(\kappa)}, \tilde{g}_j^{(\kappa)}, \Omega_{(\kappa)}$ and in Q_κ .

Note that the *first* equality is a straightforward consequence of definitions (1.8), (2.5). The *second* equality can be derived directly from the *Mehler's formula*.

Now by the *min-max* principle, for $d > 2$ and κ large enough, we obtain from the value $g_1 = \exp(-\beta/\kappa)$ the following estimates:

$$\begin{aligned} g_0 &= 1 \geq \tilde{g}_0 \geq (\Omega, \tilde{G}_\kappa(\beta)\Omega) = 1 - (\Omega, D_\kappa\Omega) \\ &= 1 - \hat{O}(\kappa^{-d/2}) > g_1 = 1 - \hat{O}(\kappa^{-1}) \geq \tilde{g}_1. \end{aligned} \quad (2.16)$$

Hence the eigenspace of the operator $\tilde{G}_\kappa(\beta)$ for its largest eigenvalue \tilde{g}_0 is *one-dimensional*. Let $\tilde{\Omega}$ be the normalized eigenfunction corresponding to \tilde{g}_0 and let $\tilde{\Omega} = a\Omega + \Omega'$, with $(\Omega, \Omega') = 0$. Then $\tilde{G}_\kappa(\beta)\tilde{\Omega} = \tilde{g}_0\tilde{\Omega}$ yields

$$a\tilde{G}_\kappa(\beta)\Omega + \tilde{G}_\kappa(\beta)\Omega' = a\tilde{g}_0\Omega + \tilde{g}_0\Omega'.$$

Applying to this relation orthogonal projector P (on Ω) and $Q = I - P$, we obtain:

$$\begin{aligned} ag_0 - a(\Omega, D_\kappa\Omega) - (\Omega, D_\kappa\Omega') &= a\tilde{g}_0, \\ -aQD_\kappa\Omega + Q\tilde{G}_\kappa(\beta)Q\Omega' &= \tilde{g}_0\Omega'. \end{aligned}$$

Since $Q\tilde{G}_\kappa(\beta)Q \leq QG_\kappa(\beta)Q \leq g_1 < \tilde{g}_0$, the operator $\tilde{g}_0 - Q\tilde{G}_\kappa(\beta)Q$ is positive and invertible. It follows from the second identity that

$$\Omega' = -a[\tilde{g}_0 - Q\tilde{G}_\kappa(\beta)Q]^{-1}QD_\kappa\Omega. \quad (2.17)$$

Together with the above first identity,

$$\begin{aligned} g_0 - \tilde{g}_0 &= (\Omega, (D_\kappa - D_\kappa Q[\tilde{g}_0 - Q\tilde{G}_\kappa(\beta)Q]^{-1}QD_\kappa)\Omega) \\ &= (W_\kappa^*\Omega, (1 - W_\kappa^*Q[\tilde{g}_0 - Q\tilde{G}_\kappa(\beta)Q]^{-1}QW_\kappa)W_\kappa^*\Omega) \end{aligned} \quad (2.18)$$

follows.

For brevity, we put

$$X' := W_\kappa^*Q[\tilde{g}_0 - QG_\kappa(\beta)Q]^{-1}QW_\kappa, \quad X := W_\kappa^*Q[1 - QG_\kappa(\beta)Q]^{-1}QW_\kappa$$

and

$$\tilde{X} := W_\kappa^*Q[\tilde{g}_0 - Q\tilde{G}_\kappa(\beta)Q]^{-1}QW_\kappa.$$

Then we get

$$\tilde{X} - X' = -\tilde{X}X',$$

and hence

$$\tilde{X} = X'(1 + X')^{-1} \quad \text{and} \quad 1 - \tilde{X} = (1 + X')^{-1}. \quad (2.19)$$

By definition of W_κ and (1.8) one gets for large κ the asymptotic:

$$W_\kappa^*\Omega = \sqrt{1 - e^{-f}}(\pi\kappa)^{-d/4}(1 + O(\kappa^{-1})).$$

By virtue of (2.18) it implies the representation:

$$g_0 - \tilde{g}_0 = (\pi\kappa)^{-d/2}(\sqrt{1 - e^{-f}}, (1 + X')^{-1}\sqrt{1 - e^{-f}})(1 + O(\kappa^{-1})) . \quad (2.20)$$

Now, we want to replace in the right hand side of this representation the operator X' by X . Note that (2.11) yields $\|X - K_f\| \leq \|X - K_f\|_1 = o(1)$. Then by $1 - \tilde{g}_0 = O(\kappa^{-d/2})$, $\tilde{g}_0 - g_1 = \hat{O}(\kappa^{-1})$, $1 - g_1 = \hat{O}(\kappa^{-1})$ and $\|W_\kappa^* \Omega\| = \hat{O}(\kappa^{-d/4})$, we find that

$$\begin{aligned} \|X' - X\| &= (1 - \tilde{g}_0)\|W_\kappa^* Q[\tilde{g}_0 - QG_\kappa(\beta)Q]^{-1}[1 - QG_\kappa(\beta)Q]^{-1}QW_\kappa\| \\ &\leq \frac{1 - \tilde{g}_0}{\tilde{g}_0 - g_1}\|W_\kappa^* Q[1 - QG_\kappa(\beta)Q]^{-1}QW_\kappa\| \\ &= O(\kappa^{1-d/2})\|X\| = O(\kappa^{1-d/2})\|K_f\| , \end{aligned}$$

which implies $\|X'\| \leq \|K_f\|(1 + o(1))$. Hence (2.20) yields

$$(\pi\kappa)^{d/2}(g_0 - \tilde{g}_0) = (\sqrt{1 - e^{-f}}, (1 + X')^{-1}\sqrt{1 - e^{-f}})(1 + O(\kappa^{-1})) \geq \frac{\|\sqrt{1 - e^{-f}}\|^2}{1 + \|K_f\|}(1 + o(1)).$$

Since the upper bound $(\pi\kappa)^{d/2}(g_0 - \tilde{g}_0) \leq (\pi\kappa)^{d/2}(\Omega, D_\kappa \Omega) \leq \|\sqrt{1 - e^{-f}}\|^2$ is obvious, we get desired estimate $g_0 - \tilde{g}_0 = \hat{O}(\kappa^{-d/2})$. Notice that the proof of the equality

$$g_0 - \tilde{g}_0 = \frac{1 + o(1)}{(\pi\kappa)^{d/2}}(\sqrt{1 - e^{-f}}, [1 + X]^{-1}\sqrt{1 - e^{-f}})$$

follows from (2.20) and the estimate:

$$\begin{aligned} &|(\sqrt{1 - e^{-f}}, (1 + X')^{-1}\sqrt{1 - e^{-f}}) - (\sqrt{1 - e^{-f}}, (1 + X)^{-1}\sqrt{1 - e^{-f}})| \\ &\leq \|\sqrt{1 - e^{-f}}\|^2\|(1 + X)^{-1}\| \|(1 + X')^{-1}\| \|X - X'\| = o(1) . \end{aligned}$$

Here $\|\sqrt{1 - e^{-f}}\|$ stands for the norm in $\mathfrak{H} = L^2(\mathbb{R}^d)$, while the other $\|\cdot\|$ for the operator norm on \mathfrak{H} . This remark finishes the proof of the lemma. \square

2.4 Evidence of Two Thermodynamic Regimes

Now we return to the conditions (2.8). We need the behavior of r_κ and \tilde{r}_κ to prove the main theorem. Here we consider the behavior of r_κ , which classifies the phase separation. That of \tilde{r}_κ is postponed to Section 4.

Proposition 2.4 (a) $\{r_\kappa\}$ converges to $r_* \in (0, 1)$ for $\kappa \rightarrow \infty$, where r_* is the unique solution of

$$\frac{\mu}{\lambda} = \frac{\log r_*}{\beta\lambda} + \frac{1}{\beta^d} \int_{[0, \infty)^d} \frac{r_* dp}{e^{|p|_1} - r_*} , \quad (2.21)$$

if and only if $\beta^d \mu < \zeta(d)\lambda$.

(b) $\kappa^d(1 - r_\kappa) \rightarrow \beta^d \lambda / (\beta^d \mu - \zeta(d)\lambda)$, and hence $\lim_{\kappa \rightarrow \infty} r_\kappa = 1$, if and only if $\beta^d \mu > \zeta(d)\lambda$.

(c) $\lim_{\kappa \rightarrow \infty} r_\kappa = 1$ and $\kappa^d(1 - r_\kappa) \rightarrow +\infty$, if and only if $\beta^d \mu = \zeta(d)\lambda$.

To this end let us introduce the notation:

$$\square_n^{(\kappa)} := \frac{\beta}{\kappa}(n + [0, 1]^d) \quad \text{for } n \in \mathbb{Z}_+^d,$$

and define for $r \in [0, 1]$, $\nu = 1, 2$ and $\kappa \in [1, \infty)$, the functions $a_\nu(\cdot; r), a_\nu^{(\kappa)}(\cdot; r)$ on $[0, \infty)^d$ by

$$a_\nu(p; r) := \frac{r e^{-|p|_1}}{(1 - r e^{-|p|_1})^\nu}$$

and by

$$a_\nu^{(\kappa)}(p; r) := \begin{cases} 0 & \text{if } p \in \square_0^{(\kappa)} \\ a_\nu(\beta n / \kappa; r) & \text{if } p \in \square_n^{(\kappa)} \end{cases} \quad \text{for } n \in \mathbb{Z}_+^d - \{0\}.$$

It is easy to show the following fact. (See [Tlb] for detail.)

Lemma 2.5 *There exists a constant c which depends only on $d > 2$ and $\beta > 0$ such that*

$$0 \leq a_\nu^{(\kappa)}(p; r) \leq a_\nu(c p; 1) \in L^1([0, \infty)^d)$$

holds for $r \in [0, 1]$ and $\nu = 1, 2$.

Remark 2.6 *If a series $\{r_\kappa\} \subset [0, 1]$ converges to $r_0 \in [0, 1]$, $a_\nu^{(\kappa)}(\cdot; r_\kappa) \rightarrow a_\nu(\cdot; r_0)$ holds a.e.. Then the lemma and the dominated convergence theorem yield*

$$\begin{aligned} \frac{\text{Tr} [r_\kappa Q_\kappa G_\kappa(\beta) Q_\kappa (1 - r_\kappa Q_\kappa G_\kappa(\beta) Q_\kappa)^{-\nu}]}{\kappa^d} &= \frac{1}{\kappa^d} \sum_{j=1}^{\infty} \frac{r_\kappa g_j^{(\kappa)}}{(1 - r_\kappa g_j^{(\kappa)})^\nu} \\ &= \left(\frac{1}{\beta^d} \right) \int_{[0, \infty)^d} a_\nu^{(\kappa)}(p; r_\kappa) dp \rightarrow \left(\frac{1}{\beta^d} \right) \int_{[0, \infty)^d} a_\nu(p; r_0) dp \end{aligned}$$

as $\kappa \rightarrow \infty$.

Proof of Proposition 2.4: If $r_\kappa \rightarrow r_* \in [0, 1)$, then by (2.8) and the above remark one gets:

$$\begin{aligned} \frac{\mu}{\lambda} &= \frac{\log r_\kappa}{\beta \lambda} + \frac{1}{\beta^d} \int_{[0, \infty)^d} a_1^{(\kappa)}(p; r_\kappa) dp + \frac{r_\kappa}{(1 - r_\kappa) \kappa^d} \\ &\longrightarrow \frac{\log r_*}{\beta \lambda} + \frac{1}{\beta^d} \int_{[0, \infty)^d} a_1(p; r_*) dp < \frac{1}{\beta^d} \int_{[0, \infty)^d} a_1(p; 1) dp = \frac{\zeta(d)}{\beta^d}. \end{aligned}$$

Since equality holds at the limit actually, (2.21) follows. Similarly, if $r_\kappa \rightarrow 1$ and $\kappa^d(1 - r_\kappa) \rightarrow \infty$, then one obtains:

$$\begin{aligned} \frac{\mu}{\lambda} &= \frac{\log r_\kappa}{\beta \lambda} + \frac{1}{\beta^d} \int_{[0, \infty)^d} a_1^{(\kappa)}(p; r_\kappa) dp + \frac{r_\kappa}{(1 - r_\kappa) \kappa^d} \\ &\longrightarrow \frac{1}{\beta^d} \int_{[0, \infty)^d} a_1(p; 1) dp = \frac{\zeta(d)}{\beta^d}. \end{aligned}$$

Finally, if $\kappa^d(1 - r_\kappa) \rightarrow \alpha > 0$, we get

$$\frac{\mu}{\lambda} = \frac{1}{\alpha} + \frac{\zeta(d)}{\beta^d} > \frac{\zeta(d)}{\beta^d} \quad \text{and} \quad \alpha = \frac{\beta^d \lambda}{\beta^d \mu - \zeta(d) \lambda}.$$

On the other hand, if $\{r_\kappa\}$ does not converge, by compactness we can take two convergent subsequences $\{r_{\kappa_i}\}$ ($i = 1, 2$) with different limits $0 < r^{(1)} < r^{(2)} < 1$. Then above arguments yield

$$\frac{\log r^{(1)}}{\beta \lambda} + \frac{1}{\beta^d} \int_{[0, \infty)^d} a_1(p; r^{(1)}) dp = \frac{\log r^{(2)}}{\beta \lambda} + \frac{1}{\beta^d} \int_{[0, \infty)^d} a_1(p; r^{(2)}) dp ,$$

which contradicts to the strict monotonicity of the function

$$h(r) = \frac{\log r}{\beta \lambda} + \frac{1}{\beta^d} \int_{[0, \infty)^d} a_1(p; r) dp . \quad (2.22)$$

Similar arguments are valid for the cases $r^{(1)} = 0$ or $r^{(2)} = 1$. If $r_\kappa \rightarrow 1$ but $\kappa^d(1 - r_\kappa)$ does not converge, we again get a contradiction. \square

3 Proof of Theorem 1.2

3.1 The case $\mu \leq \mu_c(\beta)$ (normal phase).

Let us recall that for the weak convergence of random point fields, it is enough to prove the convergence of the generating functionals. Therefore, we have to evaluate the integral in (2.4). We begin with estimates of the Fredholm determinant in the integrand. For all values of x , we have

$$\begin{aligned} & \left| \text{Det} \left(1 - (e^{ix} - 1) \frac{r_\kappa G_\kappa(\beta)}{1 - r_\kappa G_\kappa(\beta)} \right) \right|^2 \\ &= \text{Det} \left[\left(1 - (e^{ix} - 1) \frac{r_\kappa G_\kappa(\beta)}{1 - r_\kappa G_\kappa(\beta)} \right) \left(1 - (e^{-ix} - 1) \frac{r_\kappa G_\kappa(\beta)}{1 - r_\kappa G_\kappa(\beta)} \right) \right] \\ &= \text{Det} \left[1 + 4 \sin^2 \frac{x}{2} \frac{r_\kappa G_\kappa(\beta)}{(1 - r_\kappa G_\kappa(\beta))^2} \right] \geq 1. \end{aligned} \quad (3.1)$$

Set $\alpha \in (d/3, d/2)$. Then for $|x| \leq \kappa^{-\alpha}$, we have

$$\text{Det} \left[1 - (e^{ix} - 1) \frac{r_\kappa G_\kappa(\beta)}{1 - r_\kappa G_\kappa(\beta)} \right]^{-1} = \exp \left[ix \text{Tr} \frac{r_\kappa G_\kappa(\beta)}{1 - r_\kappa G_\kappa(\beta)} - \frac{x^2}{2} \text{Tr} \frac{r_\kappa G_\kappa(\beta)}{(1 - r_\kappa G_\kappa(\beta))^2} \right] (1 + O(\kappa^{d-3\alpha})) \quad (3.2)$$

In fact,

$$\log \text{ of the l.h.s.} = - \sum_{n=0}^{\infty} \log \left(1 - (e^{ix} - 1) \frac{r_\kappa g_n^{(\kappa)}}{1 - r_\kappa g_n^{(\kappa)}} \right)$$

$$= (e^{ix} - 1) \operatorname{Tr} \frac{r_\kappa G_\kappa(\beta)}{1 - r_\kappa G_\kappa(\beta)} + \frac{(e^{ix} - 1)^2}{2} \operatorname{Tr} \left(\frac{r_\kappa G_\kappa(\beta)}{1 - r_\kappa G_\kappa(\beta)} \right)^2 + R_1,$$

where $R_1 = O(\kappa^{d-3\alpha})$ since $r_\kappa \rightarrow r_* \in (0, 1)$ and

$$\operatorname{Tr} \left[\frac{r_\kappa G_\kappa(\beta)}{1 - r_\kappa G_\kappa(\beta)} \right]^\ell \leq \left(\frac{r_\kappa g_0^{(\kappa)}}{1 - r_\kappa g_0^{(\kappa)}} \right)^{\ell-1} \operatorname{Tr} \frac{r_\kappa G_\kappa(\beta)}{1 - r_\kappa G_\kappa(\beta)} = O(\kappa^d).$$

For the last equality, we recall (2.8) and Proposition 2.4(a).

Similarly, we have

$$(e^{ix} - 1) \operatorname{Tr} \frac{r_\kappa G_\kappa(\beta)}{1 - r_\kappa G_\kappa(\beta)} = \left(ix - \frac{x^2}{2} \right) \operatorname{Tr} \frac{r_\kappa G_\kappa(\beta)}{1 - r_\kappa G_\kappa(\beta)} + O(\kappa^{d-3\alpha})$$

and

$$\frac{(e^{ix} - 1)^2}{2} \operatorname{Tr} \left(\frac{r_\kappa G_\kappa(\beta)}{1 - r_\kappa G_\kappa(\beta)} \right)^2 = -\frac{x^2}{2} \operatorname{Tr} \left(\frac{r_\kappa G_\kappa(\beta)}{1 - r_\kappa G_\kappa(\beta)} \right)^2 + O(\kappa^{d-3\alpha}).$$

Thus we get (3.2) and the following lemma.

Lemma 3.1

$$\Xi_\kappa(\beta, \mu) = \frac{e^{\kappa^d(\beta\mu - \log r_\kappa)^2/2\beta\lambda} (1 + O(\kappa^{d-3\alpha}))}{\sqrt{1 + \beta\lambda\kappa^{-d} \operatorname{Tr} [r_\kappa G_\kappa(\beta)(1 - r_\kappa G_\kappa(\beta))^{-2}] \operatorname{Det}(1 - r_\kappa G_\kappa(\beta))}}$$

Proof: From the above estimates and (2.3), we have

$$\begin{aligned} & \int_{\mathbb{R}} dx \frac{e^{-is_\kappa x - \kappa^d x^2/2\beta\lambda}}{\operatorname{Det}[1 - (e^{ix} - 1)r_\kappa G_\kappa(\beta)(1 - r_\kappa G_\kappa(\beta))^{-1}]} \\ &= \int_{-\kappa^{-\alpha}}^{\kappa^{-\alpha}} dx e^{-\kappa^d x^2/2\beta\lambda - is_\kappa x + ix \operatorname{Tr} [r_\kappa G_\kappa(\beta)(1 - r_\kappa G_\kappa(\beta))^{-1}] - x^2 \operatorname{Tr} [r_\kappa G_\kappa(\beta)(1 - r_\kappa G_\kappa(\beta))^{-2}]/2} (1 + O(\kappa^{d-3\alpha})) + R_2 \\ &= \int_{-\infty}^{\infty} dx \exp \left[- \left(\frac{\kappa^d}{\beta\lambda} + \operatorname{Tr} \frac{r_\kappa G_\kappa(\beta)}{(1 - r_\kappa G_\kappa(\beta))^2} \right) \frac{x^2}{2} \right] (1 + O(\kappa^{d-3\alpha})) + R_2 + R_3, \end{aligned}$$

where

$$R_2 = \int_{|x| > \kappa^{-\alpha}} \frac{e^{-is_\kappa x - \kappa^d x^2/2\beta\lambda}}{\operatorname{Det}[1 - (e^{ix} - 1)r_\kappa G_\kappa(\beta)(1 - r_\kappa G_\kappa(\beta))^{-1}]} = O(e^{-\tilde{c}\kappa^{d-2\alpha}})$$

for some $\tilde{c} > 0$, thanks to (3.1), and

$$\begin{aligned} R_3 &= - \int_{|x| > \kappa^{-\alpha}} dx \exp \left[- \left(\frac{\kappa^d}{\beta\lambda} + \operatorname{Tr} \frac{r_\kappa G_\kappa(\beta)}{(1 - r_\kappa G_\kappa(\beta))^2} \right) \frac{x^2}{2} \right] (1 + O(\kappa^{d-3\alpha})) \\ &= O(e^{-\tilde{c}'\kappa^{d-2\alpha}}) \end{aligned}$$

for some $\tilde{c}' > 0$. Then, the lemma follows from (2.4) and (2.3). \square

For $\tilde{\Xi}_\kappa(\beta, \mu)$, we have the following asymptotics:

Lemma 3.2

$$\tilde{\Xi}_\kappa(\beta, \mu) = \frac{e^{\kappa^d(\beta\mu - \log \tilde{r}_\kappa)^2/2\beta\lambda} (1 + O(\kappa^{d-3\alpha}))}{\sqrt{1 + \beta\lambda\kappa^{-d} \text{Tr} [\tilde{r}_\kappa \tilde{G}_\kappa(\beta)(1 - \tilde{r}_\kappa \tilde{G}_\kappa(\beta))^{-2}] \text{Det}(1 - \tilde{r}_\kappa \tilde{G}_\kappa(\beta))}}$$

To show the formula, we note that "tilded" quantities are close to corresponding "untilded" ones. In fact, the following asymptotics is established. Then it is obvious to get Lemma 3.2 by a similar argument.

Lemma 3.3 *For large κ one gets:*

$$\begin{aligned} (i) \quad & \tilde{r}_\kappa \geq r_\kappa, \\ (ii) \quad & \tilde{r}_\kappa - r_\kappa = O(\kappa^{-d}), \\ (iii) \quad & \text{Tr} \left[\frac{r_\kappa G_\kappa(\beta)}{1 - r_\kappa G_\kappa(\beta)} \right] = \hat{O}(\kappa^d), \quad \text{Tr} \left[\frac{\tilde{r}_\kappa \tilde{G}_\kappa(\beta)}{1 - \tilde{r}_\kappa \tilde{G}_\kappa(\beta)} \right] = \hat{O}(\kappa^d), \\ (iv) \quad & \text{Tr} \left[\frac{r_\kappa G_\kappa(\beta)}{(1 - r_\kappa G_\kappa(\beta))^2} \right] - \text{Tr} \left[\frac{\tilde{r}_\kappa \tilde{G}_\kappa(\beta)}{(1 - \tilde{r}_\kappa \tilde{G}_\kappa(\beta))^2} \right] = O(1). \end{aligned}$$

Proof: (i) Let h_κ be the functions on $(0, 1)$ defined by

$$h_\kappa(r) = \frac{\log r}{\beta\lambda} + \frac{1}{\kappa^d} \text{Tr} \frac{r G_\kappa(\beta)}{1 - r G_\kappa(\beta)} = \frac{\log r}{\beta\lambda} + \frac{r}{\kappa^d(1 - r)} + \frac{1}{\beta^d} \int_{[0, \infty)^d} a_1^{(\kappa)}(p; r) dp.$$

We also introduce the function \tilde{h}_κ on $(0, \tilde{g}_0^{(\kappa)-1})$ by

$$\tilde{h}_\kappa(r) = \frac{\log r}{\beta\lambda} + \frac{1}{\kappa^d} \text{Tr} \frac{r \tilde{G}_\kappa(\beta)}{1 - r \tilde{G}_\kappa(\beta)}.$$

Since $G_\kappa(\beta) \geq \tilde{G}_\kappa(\beta)$, $h_\kappa \geq \tilde{h}_\kappa$ follows. Obviously h_κ and \tilde{h}_κ are strictly increasing continuous functions. Then we have $r_\kappa \leq \tilde{r}_\kappa$, because r_κ and \tilde{r}_κ are solutions of $h_\kappa(r) = \mu/\lambda$ and $\tilde{h}_\kappa(r) = \mu/\lambda$, respectively. (Recall (2.8) and (2.9).)

(ii) From $\tilde{h}_\kappa(\tilde{r}_\kappa) = h_\kappa(r_\kappa)$, we have

$$\begin{aligned} & \frac{1}{\beta\lambda} \log \frac{\tilde{r}_\kappa}{r_\kappa} \leq \tilde{h}_\kappa(\tilde{r}_\kappa) - \tilde{h}_\kappa(r_\kappa) = h_\kappa(r_\kappa) - \tilde{h}_\kappa(r_\kappa) \\ & = \frac{1}{\kappa^d} \text{Tr} \left[\frac{1}{1 - r_\kappa \tilde{G}_\kappa(\beta)} (G_\kappa(\beta) - \tilde{G}_\kappa(\beta)) \frac{r_\kappa}{1 - r_\kappa G_\kappa(\beta)} \right] = O(\kappa^{-d}). \end{aligned}$$

Here we have used Lemma 2.3 and the fact that r_κ is bounded away from 1. The desired estimate follows.

(iii) Since we already know that $r_\kappa, \tilde{r}_\kappa \rightarrow r_* \in (0, 1)$, we get these estimates readily from (2.8) and (2.9).

(iv) This can be derived by the telescoping together with (ii) and Lemma 2.3. \square

Now, let us consider the limit of the ratio $\tilde{\Xi}_\kappa(\beta, \mu)/\Xi_\kappa(\beta, \mu)$ to derive (1.27). From Lemma 3.1 and Lemma 3.2, we have

$$E_{\beta, r_*}[e^{-\langle f, \xi \rangle}] = \lim_{\kappa \rightarrow \infty} \frac{\tilde{\Xi}_\kappa(\beta, \mu)}{\Xi_\kappa(\beta, \mu)} = \lim_{\kappa \rightarrow \infty} \sqrt{\frac{1 + \beta \lambda \kappa^{-d} \text{Tr}[r_\kappa G_\kappa(\beta)(1 - r_\kappa G_\kappa(\beta))^{-2}]}{1 + \beta \lambda \kappa^{-d} \text{Tr}[\tilde{r}_\kappa \tilde{G}_\kappa(\beta)(1 - \tilde{r}_\kappa \tilde{G}_\kappa(\beta))^{-2}]}} \\ \times \frac{\text{Det}[1 - \tilde{r}_\kappa G_\kappa(\beta)] \text{Det}[1 - r_\kappa G_\kappa(\beta)]}{\text{Det}[1 - \tilde{r}_\kappa \tilde{G}_\kappa(\beta)] \text{Det}[1 - \tilde{r}_\kappa G_\kappa(\beta)]} e^{\kappa^d (2\beta\lambda)^{-1} [(\beta\mu - \log \tilde{r}_\kappa)^2 - (\beta\mu - \log r_\kappa)^2]}.$$

Lemma 3.3 yields that the first factor is equal to $1 + O(\kappa^{-d})$. For the second factor, we note that

$$\left\| \sqrt{1 - e^{-f}} \frac{\tilde{r}_\kappa G_\kappa(\beta)}{1 - \tilde{r}_\kappa G_\kappa(\beta)} \sqrt{1 - e^{-f}} - \sqrt{1 - e^{-f}} \frac{r_* G_\kappa(\beta)}{1 - r_* G_\kappa(\beta)} \sqrt{1 - e^{-f}} \right\|_1 \\ = \left\| \sqrt{1 - e^{-f}} \sqrt{\frac{r_* G_\kappa(\beta)}{1 - r_* G_\kappa(\beta)}} \frac{r_* - \tilde{r}_\kappa}{r_*} \frac{1}{1 - \tilde{r}_\kappa G_\kappa(\beta)} \sqrt{\frac{r_* G_\kappa(\beta)}{1 - r_* G_\kappa(\beta)}} \sqrt{1 - e^{-f}} \right\|_1 \\ \leq \frac{r_* - \tilde{r}_\kappa}{r_*(1 - \tilde{r}_\kappa)} \left\| \sqrt{1 - e^{-f}} \frac{r_* G_\kappa(\beta)}{1 - r_* G_\kappa(\beta)} \sqrt{1 - e^{-f}} \right\|_1 \rightarrow 0$$

holds, where $\|\cdot\|_1$ stands for the trace norm. Together with (2.10),

$$\frac{\text{Det}[1 - \tilde{r}_\kappa \tilde{G}_\kappa(\beta)]}{\text{Det}[1 - \tilde{r}_\kappa G_\kappa(\beta)]} = \text{Det}[1 + \tilde{r}_\kappa (G_\kappa(\beta) - \tilde{G}_\kappa(\beta))(1 - \tilde{r}_\kappa G_\kappa(\beta))^{-1}] \\ = \text{Det}[1 + \sqrt{1 - e^{-f}} \frac{\tilde{r}_\kappa G_\kappa(\beta)}{1 - \tilde{r}_\kappa G_\kappa(\beta)} \sqrt{1 - e^{-f}}] \rightarrow \text{Det}[1 + \sqrt{1 - e^{-f}} \frac{r_* G(\beta)}{1 - r_* G(\beta)} \sqrt{1 - e^{-f}}]$$

follows. Similarly for the third factor, we have

$$\frac{\text{Det}[1 - \tilde{r}_\kappa G_\kappa(\beta)]}{\text{Det}[1 - r_\kappa G_\kappa(\beta)]} = \text{Det}\left[1 - \frac{\tilde{r}_\kappa - r_\kappa}{r_\kappa} \frac{r_\kappa G_\kappa(\beta)}{1 - r_\kappa G_\kappa(\beta)}\right] \\ = \exp\left(-\frac{\tilde{r}_\kappa - r_\kappa}{r_\kappa} \text{Tr}\left[\frac{r_\kappa G_\kappa(\beta)}{1 - r_\kappa G_\kappa(\beta)}\right] + O(\kappa^{-d})\right) = \exp\left(-\frac{\tilde{r}_\kappa - r_\kappa}{r_\kappa} \frac{\kappa^d}{\beta\lambda} (\beta\mu - \log r_\kappa) + O(\kappa^{-d})\right),$$

where we have used Lemma 3.3 and (2.8). It also follows from Lemma 3.3 that the fourth factor is equal to

$$\exp\left[\frac{\kappa^d}{2\beta\lambda} (2\beta\mu - \log r_\kappa \tilde{r}_\kappa) \log\left(1 + \frac{r_\kappa - \tilde{r}_\kappa}{\tilde{r}_\kappa}\right)\right] = \exp\left[-\frac{\tilde{r}_\kappa - r_\kappa}{r_\kappa} \frac{\kappa^d}{\beta\lambda} (\beta\mu - \log r_\kappa) + O(\kappa^{-d})\right].$$

Thus we get (1.27). Since convergence of generating functionals yields the weak convergence of random point fields, Theorem 1.2(i) follows. \square

3.2 The case $\mu > \mu_c(\beta)$ (condensed phase).

Let us put:

$$p_j^{(\kappa)} = \frac{r_\kappa g_j^{(\kappa)}}{1 - r_\kappa g_j^{(\kappa)}}, \quad \varepsilon_j^{(\kappa)} = \log \left(1 + \frac{1}{p_j^{(\kappa)}} \right) \quad \text{and} \quad \delta^{(\kappa)} = \log \left(1 + \frac{1}{2p_1^{(\kappa)}} \right)$$

for $j = 0, 1, \dots$. Then it follows from (2.3) and Proposition 2.4(b) that

$$s_\kappa = \sum_{j=0}^{\infty} p_j^{(\kappa)}, \quad 1 + p_0^{(\kappa)} = \frac{1}{1 - r_\kappa} = \frac{\kappa^d (\beta^d \mu - \zeta(d)\lambda)}{\beta^d \lambda} (1 + o(1)) = \hat{O}(\kappa^d), \quad \varepsilon_0^{(\kappa)} = \hat{O}(\kappa^{-d}).$$

Since $g_1^{(\kappa)} = e^{-\beta/\kappa}$, we have

$$p_1^{(\kappa)} = \hat{O}(\kappa), \quad \delta^{(\kappa)} = \hat{O}(\kappa^{-1}), \quad \varepsilon_1^{(\kappa)} = \hat{O}(\kappa^{-1})$$

and

$$\varepsilon_0^{(\kappa)} < \delta^{(\kappa)} < \varepsilon_1^{(\kappa)} \leq \varepsilon_2^{(\kappa)} \leq \dots$$

Note also that $s_\kappa = \hat{O}(\kappa^d)$ holds because of (2.3), (2.8). By Remark 2.6 and $r_\kappa g_j^{(\kappa)} \in (0, 1)$

$$\sum_{j=1}^{\infty} p_j^{(\kappa)^2} = O(\kappa^d). \quad (3.3)$$

and (A.10), we get

$$\prod_{j=1}^{\infty} \left(1 - \frac{p_j^{(\kappa)}}{p_0^{(\kappa)}} \right) = \exp \left(\sum_{j=1}^{\infty} \log \left(1 - \frac{p_j^{(\kappa)}}{p_0^{(\kappa)}} \right) \right) = \exp \left(- \frac{s_\kappa - p_0^{(\kappa)}}{p_0^{(\kappa)}} + O(\kappa^{-d}) \right).$$

Similarly, we have

$$\begin{aligned} \prod_{j=1}^{\infty} (1 - p_j^{(\kappa)}(e^{\delta^{(\kappa)}} - 1)) &= \exp \left(- \sum_{j=1}^{\infty} p_j^{(\kappa)}(e^{\delta^{(\kappa)}} - 1) + O(\kappa^{d-2}) \right) \\ &= \exp \left(- \delta^{(\kappa)}(s_\kappa - p_0^{(\kappa)}) + O(\kappa^{d-2}) \right) \end{aligned}$$

since $p_j^{(\kappa)}(e^{\delta^{(\kappa)}} - 1) \leq p_1^{(\kappa)}(e^{\delta^{(\kappa)}} - 1) = 1/2$ for $j = 1, 2, \dots$.

Lemma 3.4 *For large κ we have the asymptotics:*

$$\Xi_\kappa(\beta, \mu) = \sqrt{\frac{2\pi\beta\lambda}{e^2\kappa^d}} \frac{\beta^{d-1} e^{\kappa^d(\beta\mu - \log r_\kappa)^2/2\beta\lambda}}{(\beta^d\mu - \zeta(d)\lambda)\text{Det}(1 - r_\kappa G_\kappa(\beta))} (1 + o(1)). \quad (3.4)$$

Proof: As in the proof of Lemma 3.1, we start with the integral

$$\mathcal{I} = \int_{\mathbb{R}} dx \frac{e^{-is_\kappa x - \kappa^d x^2/2\beta\lambda}}{\text{Det}[1 - (e^{ix} - 1)r_\kappa G_\kappa(\beta)(1 - r_\kappa G_\kappa(\beta))^{-1}]}$$

$$= \int_{-\infty}^{\infty} dz \frac{e^{-is_{\kappa}z - \kappa^d z^2/2\beta\lambda}}{\prod_{j=0}^{\infty} (1 - (e^{iz} - 1)p_j^{(\kappa)})} = \left(\int_{-\infty}^{\infty} - \int_{-\infty - i\delta^{(\kappa)}}^{\infty - i\delta^{(\kappa)}} \right) + \int_{-\infty - i\delta^{(\kappa)}}^{\infty - i\delta^{(\kappa)}} = \mathcal{I}_1 + \mathcal{I}_2.$$

Since $1 + p_0^{(\kappa)} - e^{iz}p_0^{(\kappa)} = (1 + p_0^{(\kappa)})(1 - e^{i(z - 2n\pi + i\epsilon_0^{(\kappa)})})$, the integrand of \mathcal{I}_1 has simple poles at $z = 2n\pi - i\epsilon_0^{(\kappa)}$ ($n \in \mathbb{Z}$). Let us calculate \mathcal{I}_1 by means of residues:

$$\begin{aligned} \mathcal{I}_1 &= -2\pi i \sum_{n=-\infty}^{\infty} \text{Res} \left[\frac{1}{(1 + p_0^{(\kappa)})(1 - e^{i(z - 2n\pi + i\epsilon_0^{(\kappa)})})} \frac{e^{-is_{\kappa}z - \kappa^d z^2/2\beta\lambda}}{\prod_{j=1}^{\infty} (1 - (e^{iz} - 1)p_j^{(\kappa)})}; 2n\pi - i\epsilon_0^{(\kappa)} \right] \\ &= \frac{2\pi e^{-s_{\kappa}\epsilon_0^{(\kappa)} + \kappa^d \epsilon_0^{(\kappa)2}/2\beta\lambda}}{(1 + p_0^{(\kappa)}) \prod_{j=1}^{\infty} (1 - p_j^{(\kappa)}/p_0^{(\kappa)})} (1 + O(e^{-2\pi^2 \kappa^d/\beta\lambda})) \\ &= \frac{2\pi \beta^d \lambda e^{-1 + O(\kappa^{-d})}}{\kappa^d (\beta^d \mu - \zeta(d)\lambda)}. \end{aligned}$$

Here the pole $z = -i\epsilon_0^{(\kappa)}$ gives the dominant contribution in the second equality. In the third equality, we have used the relations above this Lemma.

On the other hand, we have

$$\begin{aligned} |\mathcal{I}_2| &\leq \int_{-\infty}^{\infty} dx \frac{e^{-s_{\kappa}\delta^{(\kappa)} - \kappa^d(x^2 - \delta^{(\kappa)2})/2\beta\lambda}}{\prod_{j=0}^{\infty} |(1 - (e^{ix + \delta^{(\kappa)}} - 1)p_j^{(\kappa)})|} \\ &\leq \sqrt{\frac{2\pi\beta\lambda}{\kappa^d}} \frac{e^{-\hat{O}(\kappa^{d-1})}}{(e^{\delta^{(\kappa)}} - 1)p_0^{(\kappa)} - 1} = o(\mathcal{I}_1), \end{aligned}$$

where we have used

$$\begin{aligned} \prod_{j=0}^{\infty} |(1 - (e^{ix + \delta^{(\kappa)}} - 1)p_j^{(\kappa)})| &\geq ((e^{\delta^{(\kappa)}} - 1)p_0^{(\kappa)} - 1) \prod_{j=1}^{\infty} |(1 - (e^{\delta^{(\kappa)}} - 1)p_j^{(\kappa)})| \\ &\geq ((e^{\delta^{(\kappa)}} - 1)p_0^{(\kappa)} - 1) \exp(-\delta^{(\kappa)}(s_{\kappa} - p_0^{(\kappa)}) + O(\kappa^{d-2})), \end{aligned}$$

which follows from

$$1 - (e^{ix + \delta^{(\kappa)}} - 1)p_j^{(\kappa)} = (1 + p_j^{(\kappa)})(1 - e^{ix + \delta^{(\kappa)} - \varepsilon_j^{(\kappa)}})$$

and $\delta^{(\kappa)} - \varepsilon_0^{(\kappa)} > 0$, $\delta^{(\kappa)} - \varepsilon_j^{(\kappa)} < 0$ ($j = 1, 2, \dots$). Note also that $\delta^{(\kappa)}p_0^{(\kappa)} = \hat{O}(\kappa^{d-1})$ holds. From (2.4) and the first equality in (2.3) one gets desired expression for the asymptotics of $\Xi_{\kappa}(\beta, \mu)$. \square

In order to obtain the corresponding asymptotics for $\tilde{\Xi}_{\kappa}(\beta, \mu)$, we use the following estimates about $\tilde{p}_j^{(\kappa)} = \tilde{r}_{\kappa}\tilde{g}_j^{(\kappa)}/(1 - \tilde{r}_{\kappa}\tilde{g}_j^{(\kappa)})$ ($j = 0, 1, \dots$).

Lemma 3.5

$$\begin{aligned} 1 - \tilde{r}_{\kappa}\tilde{g}_0^{(\kappa)} &= \frac{\beta^d \lambda (1 + o(1))}{\kappa^d (\beta^d \mu - \zeta(d)\lambda)}, \quad |1 - \tilde{r}_{\kappa}| = O(\kappa^{-d/2}), \\ \tilde{p}_0^{(\kappa)} &= \hat{O}(\kappa^d), \quad \tilde{p}_1^{(\kappa)} = \hat{O}(\kappa), \quad \sum_{j=1}^{\infty} \tilde{p}_j^{(\kappa)} = \hat{O}(\kappa^d), \quad \sum_{j=1}^{\infty} \tilde{p}_j^{(\kappa)2} = O(\kappa^d). \end{aligned}$$

Proof : Proposition 2.4(b) , Lemma 3.3(i) and Lemma 2.3 yield

$$1 - \hat{O}(\kappa^{-d}) = r_\kappa \leq \tilde{r}_\kappa < \tilde{g}_0^{(\kappa)^{-1}} = 1 + \hat{O}(\kappa^{-d/2}), \quad (3.5)$$

which implies $|1 - \tilde{r}_\kappa| = O(\kappa^{-d/2})$. Note that the argument which shows $r_\kappa \leq \tilde{r}_\kappa$ in the proof of Lemma 3.3 is also valid for the present case. In the variational formula

$$\tilde{g}_1^{(\kappa)} = \sup_{\psi \perp \tilde{\Omega}} \frac{(\psi, \tilde{G}_\kappa(\beta)\psi)}{(\psi, \psi)} = \sup_{\psi \perp \tilde{\Omega}} \frac{[(\psi, G_\kappa(\beta)\psi) - (\psi, D_\kappa\psi)]}{(\psi, \psi)},$$

we can use as ψ a linear combination of two eigenfunctions $\kappa^{-d/4}\phi_s(x/\sqrt{\kappa})$ of $G_\kappa(\beta)$ perpendicular to $\tilde{\Omega}$ (e.g., with $s = (1, 0, 0, \dots)$ and $s = (0, 1, 0, \dots)$). Then we get $\tilde{g}_1^{(\kappa)} \geq 1 - \hat{O}(\kappa^{-1})$. Together with $\tilde{g}_1^{(\kappa)} \leq g_1^{(\kappa)} = 1 - \hat{O}(\kappa^{-1})$, $\tilde{g}_1^{(\kappa)} = 1 - \hat{O}(\kappa^{-1})$ follows. Thus $\hat{O}(\kappa) = \tilde{p}_1^{(\kappa)} \geq \tilde{p}_2^{(\kappa)} \geq \dots$ holds.

Now we get

$$\begin{aligned} \left| \sum_{j=1}^{\infty} \tilde{p}_j^{(\kappa)} - \sum_{j=1}^{\infty} \tilde{g}_j^{(\kappa)} / (1 - \tilde{g}_j^{(\kappa)}) \right| &= |1 - \tilde{r}_\kappa| \sum_{j=1}^{\infty} \tilde{g}_j^{(\kappa)} / ((1 - \tilde{r}_\kappa \tilde{g}_j^{(\kappa)})(1 - \tilde{g}_j^{(\kappa)})) \\ &= O(\kappa^{1-d/2}) \sum_{j=1}^{\infty} \tilde{g}_j^{(\kappa)} / (1 - \tilde{g}_j^{(\kappa)}), \end{aligned}$$

which implies

$$\sum_{j=1}^{\infty} \tilde{p}_j^{(\kappa)} = (1 + O(\kappa^{1-d/2})) \sum_{j=1}^{\infty} \tilde{g}_j^{(\kappa)} / (1 - \tilde{g}_j^{(\kappa)}).$$

On the other hand, because

$$\left| \sum_{j=1}^{\infty} \frac{\tilde{g}_j^{(\kappa)}}{1 - \tilde{g}_j^{(\kappa)}} - \sum_{j=1}^{\infty} \frac{g_j^{(\kappa)}}{1 - g_j^{(\kappa)}} \right| \leq \sum_{j=1}^{\infty} \frac{g_j^{(\kappa)} - \tilde{g}_j^{(\kappa)}}{(1 - g_1^{(\kappa)})(1 - \tilde{g}_1^{(\kappa)})} \leq \frac{\text{Tr } D_\kappa}{(1 - g_1^{(\kappa)})(1 - \tilde{g}_1^{(\kappa)})} = O(\kappa^2),$$

we have

$$\frac{1}{\kappa^d} \sum_{j=1}^{\infty} \frac{\tilde{g}_j^{(\kappa)}}{1 - \tilde{g}_j^{(\kappa)}} = \frac{1}{\kappa^d} \sum_{j=1}^{\infty} \frac{g_j^{(\kappa)}}{1 - g_j^{(\kappa)}} + O(\kappa^{2-d}) = \frac{\zeta(d)}{\beta^d} + o(1),$$

where we recall Remark 2.6. Thus we have $\kappa^{-d} \sum_{j=1}^{\infty} \tilde{p}_j^{(\kappa)} = \beta^{-d} \zeta(d) + o(1)$. Using (2.9), we get

$$\frac{\tilde{r}_\kappa \tilde{g}_0^{(\kappa)}}{\kappa^d (1 - \tilde{r}_\kappa \tilde{g}_0^{(\kappa)})} = -\frac{\log \tilde{r}_\kappa}{\beta \lambda} + \frac{\mu}{\lambda} - \frac{1}{\kappa^d} \sum_{j=1}^{\infty} \tilde{p}_j^{(\kappa)} = \frac{\mu}{\lambda} - \frac{\zeta(d)}{\beta^d} + o(1)$$

which yields the first and the third equality.

To prove the remaining last bound, it is enough to show that

$$\tilde{p}_j^{(\kappa)} \leq 2p_j^{(\kappa)} \quad (j = 1, 2, \dots) \quad (3.6)$$

hold for large enough κ , because of $\sum_{j=1}^{\infty} p_j^{(\kappa)^2} = O(\kappa^d)$. In fact, in the expression

$$\tilde{p}_j^{(\kappa)} = \frac{r_\kappa \tilde{g}_j^{(\kappa)}}{1 - r_\kappa \tilde{g}_j^{(\kappa)}} \frac{1 + (\tilde{r}_\kappa - r_\kappa)/r_\kappa}{1 - (\tilde{r}_\kappa - r_\kappa) \tilde{g}_j^{(\kappa)} / (1 - r_\kappa \tilde{g}_j^{(\kappa)})},$$

$(\tilde{r}_\kappa - r_\kappa)/r_\kappa = O(\kappa^{-d/2})$ and $|(\tilde{r}_\kappa - r_\kappa) \tilde{g}_j^{(\kappa)} / (1 - r_\kappa \tilde{g}_j^{(\kappa)})| \leq (\tilde{r}_\kappa - r_\kappa) / (1 - r_\kappa \tilde{g}_1^{(\kappa)}) = O(\kappa^{1-d/2})$ hold. Because of $\tilde{g}_j^{(\kappa)} \leq g_j^{(\kappa)}$, we also have $r_\kappa \tilde{g}_j^{(\kappa)} / (1 - r_\kappa \tilde{g}_j^{(\kappa)}) \leq p_j^{(\kappa)}$. Thus we get (3.6). \square

It is obvious now that the next Lemma can be derived along the same line of reasoning as the proof of Lemma 3.4.

Lemma 3.6 *For large κ one gets the asymptotics:*

$$\tilde{\Xi}_\kappa(\beta, \mu) = \sqrt{\frac{2\pi\beta\mu}{e^2\kappa^d}} \frac{\beta^{d-1} e^{\kappa^d(\beta\mu - \log \tilde{r}_\kappa)^2 / 2\beta\lambda}}{(\beta^d\mu - \zeta(d)\lambda) \text{Det}(1 - \tilde{r}_\kappa \tilde{G}_\kappa(\beta))} (1 + o(1)). \quad (3.7)$$

In order to calculate the limit of $\tilde{\Xi}_\kappa(\beta, \mu) / \Xi_\kappa(\beta, \mu)$, we use the following lemma, where we put

$$\hat{g}_0^{(\kappa)} := (\Omega_\kappa, \tilde{G}_\kappa(\beta) \Omega_\kappa) + \tilde{r}_\kappa (\Omega_\kappa, \tilde{G}_\kappa(\beta) Q_\kappa (1 - \tilde{r}_\kappa Q_\kappa \tilde{G}_\kappa(\beta) Q_\kappa)^{-1} Q_\kappa \tilde{G}_\kappa(\beta) \Omega_\kappa).$$

Lemma 3.7 *For large κ one gets:*

$$\begin{aligned} \text{(i)} \quad & \tilde{r}_\kappa - r_\kappa = (1 - \tilde{g}_0^{(\kappa)})(1 + o(1)) = \hat{O}(\kappa^{-d/2}), \\ \text{(ii)} \quad & 1 - \tilde{r}_\kappa \hat{g}_0^{(\kappa)} = (1 - \tilde{r}_\kappa \tilde{g}_0^{(\kappa)})(1 + o(1)). \end{aligned}$$

Proof: From Lemma 3.5 and Proposition 2.4(b), we have $\tilde{r}_\kappa \tilde{g}_0^{(\kappa)} - r_\kappa = o(\kappa^{-d})$. Hence (i) follows from $\tilde{r}_\kappa - r_\kappa = \tilde{r}_\kappa(1 - \tilde{g}_0^{(\kappa)}) + \tilde{r}_\kappa \tilde{g}_0^{(\kappa)} - r_\kappa$, $\tilde{r}_\kappa = 1 + O(\kappa^{d/2})$ and Lemma 2.3.

By virtue of Lemma 2.3 we get

$$\tilde{g}_0^{(\kappa)} - \hat{g}_0^{(\kappa)} = (W_\kappa^* \Omega_\kappa, W_\kappa^* Q_\kappa [(\tilde{g}_0^{(\kappa)} - Q_\kappa \tilde{G}_\kappa(\beta) Q_\kappa)^{-1} - (\tilde{r}_\kappa^{-1} - Q_\kappa \tilde{G}_\kappa(\beta) Q_\kappa)^{-1}] Q_\kappa W_\kappa W_\kappa^* \Omega_\kappa)$$

$$= (W_\kappa^* \Omega_\kappa, W_\kappa^* Q_\kappa (\tilde{g}_0^{(\kappa)} - Q_\kappa \tilde{G}_\kappa(\beta) Q_\kappa)^{-1} (\tilde{r}_\kappa^{-1} - \tilde{g}_0^{(\kappa)}) (\tilde{r}_\kappa^{-1} - Q_\kappa \tilde{G}_\kappa(\beta) Q_\kappa)^{-1} Q_\kappa W_\kappa W_\kappa^* \Omega_\kappa), \quad (3.8)$$

which yields $\tilde{g}_0^{(\kappa)} - \hat{g}_0^{(\kappa)} \geq 0$ since $\tilde{r}_\kappa < \tilde{g}_0^{(\kappa)-1}$. Recall that $\|\tilde{X}\| \leq 1$ and $\|W_\kappa^* \Omega_0^{(\kappa)}\| = O(\kappa^{-d/4})$. (See (2.19) and its next line.) Then we also get from (3.8) that

$$\begin{aligned} \tilde{g}_0^{(\kappa)} - \hat{g}_0^{(\kappa)} &\leq \frac{1 - \tilde{r}_\kappa \tilde{g}_0^{(\kappa)}}{1 - \tilde{r}_\kappa \tilde{g}_1^{(\kappa)}} (W_\kappa^* \Omega_0^{(\kappa)}, W_\kappa^* Q_\kappa (\tilde{g}_0^{(\kappa)} - Q_\kappa \tilde{G}_\kappa(\beta) Q_\kappa)^{-1} Q_\kappa W_\kappa W_\kappa^* \Omega_0^{(\kappa)}) \\ &\leq \frac{1 - \tilde{r}_\kappa \tilde{g}_0^{(\kappa)}}{1 - \tilde{r}_\kappa \tilde{g}_1^{(\kappa)}} \|\tilde{X}\| \|W_\kappa^* \Omega_0^{(\kappa)}\|^2 = (1 - \tilde{r}_\kappa \tilde{g}_0^{(\kappa)}) O(\kappa^{1-d/2}). \end{aligned}$$

Hence we obtain the asymptotics (ii):

$$1 - \tilde{r}_\kappa \hat{g}_0^{(\kappa)} = 1 - \tilde{r}_\kappa \tilde{g}_0^{(\kappa)} + \tilde{r}_\kappa (\tilde{g}_0^{(\kappa)} - \hat{g}_0^{(\kappa)}) = (1 - \tilde{r}_\kappa \tilde{g}_0^{(\kappa)})(1 + o(1)). \quad \square$$

Now, taking into account (3.4) and (3.7), we can find the asymptotics of the generating functional (1.25):

$$\begin{aligned} E_{\kappa, \beta, \mu} [e^{-\langle f, \xi \rangle}] &= \frac{\tilde{\Xi}_\kappa(\beta, \mu)}{\Xi_\kappa(\beta, \mu)} = \exp \left(\frac{\kappa^d}{2\beta\lambda} (2\beta\mu - \log r_\kappa \tilde{r}_\kappa) \log \frac{r_\kappa}{\tilde{r}_\kappa} \right) \frac{\text{Det}[1 - \tilde{r}_\kappa Q_\kappa \tilde{G}_\kappa(\beta) Q_\kappa]}{\text{Det}[1 - \tilde{r}_\kappa \tilde{G}_\kappa(\beta)]} \\ &\times \frac{\text{Det}[1 - \tilde{r}_\kappa Q_\kappa G_\kappa(\beta) Q_\kappa]}{\text{Det}[1 - \tilde{r}_\kappa Q_\kappa \tilde{G}_\kappa(\beta) Q_\kappa]} \frac{\text{Det}[1 - r_\kappa Q_\kappa G_\kappa(\beta) Q_\kappa]}{\text{Det}[1 - \tilde{r}_\kappa Q_\kappa G_\kappa(\beta) Q_\kappa]} \frac{\text{Det}[1 - r_\kappa G_\kappa(\beta)]}{\text{Det}[1 - r_\kappa Q_\kappa G_\kappa(\beta) Q_\kappa]} (1 + o(1)). \end{aligned} \quad (3.9)$$

By virtue of Lemma 3.7(i), for the exponent of the first factor, we have

$$\frac{\kappa^d}{2\beta\lambda} (2\beta\mu - \log r_\kappa \tilde{r}_\kappa) \log \frac{r_\kappa}{\tilde{r}_\kappa} = -\frac{\mu \kappa^d}{\lambda} (1 - \tilde{g}_0^{(\kappa)})(1 + o(1)).$$

For the second factor, we use the Feshbach formula, which claims

$$\text{Det} A = \text{Det} B \text{Det}(C - K^T B^{-1} K),$$

where

$$A = \begin{pmatrix} B & -K \\ -K^T & C \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ -K^T B^{-1} & 1 \end{pmatrix} \begin{pmatrix} B & 0 \\ 0 & C - K^T B^{-1} K \end{pmatrix} \begin{pmatrix} 1 & -B^{-1} K \\ 0 & 1 \end{pmatrix}.$$

This formula and Lemma 3.7(ii) yield

$$\begin{aligned} \frac{\text{Det}[1 - \tilde{r}_\kappa Q_\kappa \tilde{G}_\kappa(\beta) Q_\kappa]}{\text{Det}[1 - \tilde{r}_\kappa \tilde{G}_\kappa(\beta)]} &= \frac{1}{1 - \tilde{r}_\kappa (\Omega_0^{(\kappa)}, \tilde{G}_\kappa(\beta) \Omega_0^{(\kappa)}) - (\Omega_0^{(\kappa)}, \tilde{r}_\kappa \tilde{G}_\kappa(\beta) Q_\kappa (1 - \tilde{r}_\kappa Q_\kappa \tilde{G}_\kappa(\beta) Q_\kappa)^{-1} Q_\kappa \tilde{r}_\kappa \tilde{G}_\kappa(\beta) \Omega_0^{(\kappa)})} \\ &= 1 / (1 - \tilde{r}_\kappa \hat{g}_0^{(\kappa)}) = (1 + o(1)) / (1 - \tilde{r}_\kappa \tilde{g}_0^{(\kappa)}). \end{aligned}$$

Since

$$\frac{\text{Det}[1 - r_\kappa G_\kappa(\beta)]}{\text{Det}[1 - r_\kappa Q_\kappa G_\kappa(\beta) Q_\kappa]} = 1 - r_\kappa,$$

then Proposition 2.4(b) and Lemma 3.5 yields for the product of factors in (3.9):

$$(\text{the 2nd factor}) \times (\text{the last factor}) \rightarrow 1$$

in the limit $\kappa \rightarrow \infty$.

Now, since Lemma 3.5 and (2.11) give

$$\| \sqrt{1 - e^{-f}} \frac{\tilde{r}_\kappa Q_\kappa G_\kappa(\beta) Q_\kappa}{1 - \tilde{r}_\kappa Q_\kappa G_\kappa(\beta) Q_\kappa} \sqrt{1 - e^{-f}} - K_f \|_1$$

$$\begin{aligned}
&\leq \left\| \sqrt{1-e^{-f}} \frac{\tilde{r}_\kappa Q_\kappa G_\kappa(\beta) Q_\kappa}{1 - \tilde{r}_\kappa Q_\kappa G_\kappa(\beta) Q_\kappa} \sqrt{1-e^{-f}} - \sqrt{1-e^{-f}} \frac{Q_\kappa G_\kappa(\beta) Q_\kappa}{1 - Q_\kappa G_\kappa(\beta) Q_\kappa} \sqrt{1-e^{-f}} \right\|_1 \\
&\quad + \left\| \sqrt{1-e^{-f}} \frac{Q_\kappa G_\kappa(\beta) Q_\kappa}{1 - Q_\kappa G_\kappa(\beta) Q_\kappa} \sqrt{1-e^{-f}} - K_f \right\|_1 \\
&\leq |\tilde{r}_\kappa - 1| \left\| \sqrt{1-e^{-f}} \frac{Q_\kappa G_\kappa(\beta) Q_\kappa}{1 - Q_\kappa G_\kappa(\beta) Q_\kappa} \sqrt{1-e^{-f}} \right\|_1 \|(1 - \tilde{r}_\kappa Q_\kappa G_\kappa(\beta) Q_\kappa)^{-1}\| \\
&\quad + \left\| \sqrt{1-e^{-f}} \frac{Q_\kappa G_\kappa(\beta) Q_\kappa}{1 - Q_\kappa G_\kappa(\beta) Q_\kappa} \sqrt{1-e^{-f}} - K_f \right\|_1 \rightarrow 0
\end{aligned}$$

for $\kappa \rightarrow \infty$, we obtain the limit:

$$\begin{aligned}
&\frac{\text{Det}[1 - \tilde{r}_\kappa Q_\kappa G_\kappa(\beta) Q_\kappa]}{\text{Det}[1 - \tilde{r}_\kappa Q_\kappa \tilde{G}_\kappa(\beta) Q_\kappa]} = \frac{1}{\text{Det}[1 + \tilde{r}_\kappa Q_\kappa (G_\kappa(\beta) - \tilde{G}_\kappa(\beta)) Q_\kappa (1 - \tilde{r}_\kappa Q_\kappa G_\kappa(\beta) Q_\kappa)^{-1}]} \\
&= \text{Det}[1 + \tilde{r}_\kappa \sqrt{1-e^{-f}} Q_\kappa G_\kappa(\beta) Q_\kappa (1 - \tilde{r}_\kappa Q_\kappa G_\kappa(\beta) Q_\kappa)^{-1} \sqrt{1-e^{-f}}]^{-1} \rightarrow \text{Det}[1 + K_f]^{-1}
\end{aligned}$$

for the third factor in (3.9). Here we have used the *cyclicity* of the Fredholm determinant.

Lemma 3.7(i), (3.3), (2.3) and Proposition 2.4(b) yield

$$\begin{aligned}
&\frac{\text{Det}[1 - r_\kappa Q_\kappa G_\kappa(\beta) Q_\kappa]}{\text{Det}[1 - \tilde{r}_\kappa Q_\kappa G_\kappa(\beta) Q_\kappa]} = \frac{1}{\text{Det}[1 - (\tilde{r}_\kappa - r_\kappa) Q_\kappa G_\kappa(\beta) Q_\kappa (1 - r_\kappa Q_\kappa G_\kappa(\beta) Q_\kappa)^{-1}]} \\
&= \exp \left(\frac{\tilde{r}_\kappa - r_\kappa}{r_\kappa} \text{Tr} \frac{r_\kappa Q_\kappa G_\kappa(\beta) Q_\kappa}{1 - r_\kappa Q_\kappa G_\kappa(\beta) Q_\kappa} + O(1) \right) = \exp \left((1 - \tilde{g}_0^{(\kappa)}) \left[\kappa^d \frac{\beta \mu - \log r_\kappa}{\beta \lambda} - \frac{r_\kappa}{1 - r_\kappa} \right] + O(1) \right) \\
&= \exp \{ (1 - \tilde{g}_0^{(\kappa)}) \kappa^d \zeta(d) (1 + o(1)) / \beta^d \}.
\end{aligned}$$

Thus, by Lemma 2.3 and (2.11) we get for the product in (3.9):

$$\begin{aligned}
&(\text{the 1st factor}) \times (\text{the 4th factor}) \\
&= \exp \left(- \frac{\kappa^{d/2} (1 + o(1))}{\pi^{d/2}} \frac{\beta^d \mu - \zeta(d) \lambda}{\beta^d \lambda} (\sqrt{1-e^{-f}}, (1 + K_f)^{-1} \sqrt{1-e^{-f}}) \right).
\end{aligned}$$

Now Theorem 1.2(ii) follows by collecting in (3.9) the asymptotics of factors that we find above. \square

4 Proof of Theorem 1.8

We start with the grand-canonical expectation value of the total number of MF interacting bosons in the WHT (1.31):

$$N_{\kappa, \lambda}(\beta, \mu) := \kappa^d \rho_{\kappa, \lambda}^{(tot)}(\beta, \mu) = \frac{1}{\beta \Xi_\kappa(\beta, \mu)} \frac{\partial}{\partial \mu} \Xi_\kappa(\beta, \mu),$$

where we use for $\Xi_\kappa(\beta, \mu)$ the expression (2.2) after the z -integration where the values of s and r are not fixed yet. The differentiation with respect to μ can be converted

into differentiation with respect to x in the Fredholm determinant. Then integrating by parts we obtain

$$N_{\kappa,\lambda}(\beta, \mu) = s - \frac{i\kappa^d}{\beta\lambda} R(r, s) ,$$

where

$$\begin{aligned} R(r, s) &:= \int_{\mathbb{R}} dx \frac{x e^{-isx - \kappa^d x^2 / 2\beta\lambda}}{\text{Det}[1 - (e^{ix} - 1)rG_{\kappa}(\beta)(1 - rG_{\kappa}(\beta))^{-1}]} \\ &\times \left[\int_{\mathbb{R}} dx \frac{e^{-isx - \kappa^d x^2 / 2\beta\lambda}}{\text{Det}[1 - (e^{ix} - 1)rG_{\kappa}(\beta)(1 - rG_{\kappa}(\beta))^{-1}]} \right]^{-1} . \end{aligned} \quad (4.1)$$

Then we put $s = s_{\kappa}$ and $r = r_{\kappa}$ such that (2.3) holds, and if we prove $R(r_{\kappa}, s_{\kappa}) = o(1)$, then (1.32) follows as a consequence of this asymptotics.

To this end notice that for the case (i) $\mu < \mu_c(\beta)$, we get $R(r_{\kappa}, s_{\kappa}) = O(\kappa^{-d/2})$ from the estimates in the proof of Lemma 3.1.

For the case (ii) $\mu > \mu_c(\beta)$, using the notations of the proof of Lemma 3.4, one obtains for the first factor in (4.1):

$$\int_{\mathbb{R}} dx \frac{x e^{-is_{\kappa}x - \kappa^d x^2 / 2\beta\lambda}}{\text{Det}[1 - (e^{ix} - 1)r_{\kappa}G_{\kappa}(\beta)(1 - r_{\kappa}G_{\kappa}(\beta))^{-1}]} = -i\varepsilon_0 \mathcal{I}_1 + o(\mathcal{I}_1) .$$

Therefore, we get that $R(r_{\kappa}, s_{\kappa}) = O(\kappa^{-d})$.

The other properties stated in the Theorem 1.8 follow straightforwardly from Section 2.4 and the line of reasoning developed for the proof of Proposition 2.4. \square

5 Concluding Remarks and Conjectures

In the present paper we consider a model of mean-field interacting boson gas in traps described by the harmonic potential. For this model we study the position distribution of the constituent bosons in the WHT limit by means of the RPF method.

It is shown that there are two phases distinguished by the boson condensation. In one domain of parameters, the resulting generating functional for the RPF is the same as for the non-interacting boson gas, for unconventional values of the IBG parameters. Whereas in another domain, the generating functional describes divergence of the density due to the localization of macroscopic number of particles.

Our results are obtained via analysis of the generating functional. We do not intend to start the analysis using the characteristic functional here. However, we would like to mention a topic on the central limit theorem as a conjecture.

Let us consider $E_{\kappa,\beta,\mu}[e^{-\langle f, \xi \rangle}]$ for small $f \in C_0(\mathbb{R}^d)$ in the sense of the sup-norm $\|\cdot\|_{\infty}$. By Theorem 1.2(ii) we obtain:

$$\begin{aligned} E_{\kappa,\beta,\mu}[e^{-\langle f, \xi \rangle}] &= \\ \exp \left\{ -\frac{\kappa^{d/2}}{\pi^{d/2}} \frac{\mu - \mu_{\lambda,c}(\beta)}{\lambda} \left[\int_{\mathbb{R}^d} dx f(x) + \frac{1}{2} \langle f, (1 + G(\beta))(1 - G(\beta))^{-1} f \rangle + \dots \right] \right\} . \end{aligned}$$

This leads us to conclusion that the following claim might be true:

Conjecture 5.1 *Let ξ_κ be the random point measure on \mathbb{R}^d with distribution given by $\nu_{\kappa,\beta,\mu}$ for $\kappa > 0$, and \mathfrak{L} be the Lebesgue measure on \mathbb{R}^d . Then for $\kappa \rightarrow \infty$ the random field*

$$\frac{\pi^{d/4}}{\kappa^{d/4}} \left(\frac{\lambda}{\mu - \mu_{\lambda,c}(\beta)} \right)^{1/2} \left(\xi_\kappa - \frac{\kappa^{d/2}}{\pi^{d/2}} \frac{\mu - \mu_{\lambda,c}(\beta)}{\lambda} \mathfrak{L} \right)$$

converges in distribution to the Gaussian random field on \mathbb{R}^d with covariance

$$(1 + G(\beta))(1 - G(\beta))^{-1} .$$

We finish by some remarks about the method of the RPF approach to the BEC in the WHT limit used in the present paper:

- (i) It could be applied to a general “non-quadratic” mean-field interaction $U_\Phi := \kappa^d \Phi(n/\kappa^d)$, where $\Phi : x \in \mathbb{R} \mapsto \mathbb{R}$ is a piece-wise differentiable continuous function bounded from below [TZ], as well as to the van der Waals particle interaction, which is more local than the mean-field [deS-Z]. We guess that for this kind of interaction the particle distribution will spread as κ^α with some large α even for condensed particles.
- (ii) The method has to be compared with the scaled external field perturbation of BEC considered in [deS-Z], [Pu]. We suppose that this could clarify the concept of the choice of “effective” volume, since it is important for description of the local particle density measured in the WHT limit BEC experiments as well as its definition of the mean-field interaction [DGPS], [LSSY], [PeSm].

Acknowledgements

H.T. thanks MEXT for the financial support under the Grant-in-Aid for Scientific Research No. 17654021. V.A.Z. is thankful to Mathematical Department of the Kanazawa University for a warm hospitality and financial support.

A Miscellaneous formulae

$$\frac{x}{1 - e^{-x}} \leq 1 + x \leq 2 \vee (2x) \quad \text{for} \quad x > 0. \quad (\text{A.1})$$

$$0 \leq \left(\frac{2x}{1 - e^{-2x}} \right)^{d/2} - 1 \leq Ax \quad \text{for} \quad x \in [0, 1], \quad (\text{A.2})$$

$$0 \leq \left(\frac{1}{1 - e^{-2x}} \right)^{d/2} - 1 \leq Be^{-2x} \quad \text{for} \quad x \geq 1, \quad (\text{A.3})$$

where A and B are constants depending only on d .

$$1 - e^{-x} \leq x \quad \text{for} \quad x \geq 0. \quad (\text{A.4})$$

$$e^{-x} - e^{-y} \leq \frac{1}{e} \left(\frac{y}{x} - 1 \right) \quad \text{for} \quad 0 < x \leq y. \quad (\text{A.5})$$

$$\tanh x \leq x \wedge 1 \quad \text{for} \quad x \geq 0. \quad (\text{A.6})$$

$$|\coth x - 1| \leq \frac{2e}{e - 1} e^{-x} \quad \text{for} \quad x \geq 1. \quad (\text{A.7})$$

$$1 \leq \frac{\sinh x}{x} \leq 1 + x^2 \quad \text{for} \quad x \in [0, 1]. \quad (\text{A.8})$$

$$\frac{1}{\sinh x} \leq \frac{2}{1 - 1/e^2} e^{-x} \quad \text{for} \quad x \geq 1. \quad (\text{A.9})$$

$$| -\log(1 - x) - x | \leq \frac{x^2}{2(1 - x)} \quad \text{for} \quad x < 1. \quad (\text{A.10})$$

References

- [BrRo] O. Bratteli and D. W. Robinson, *Operator algebras and quantum statistical mechanics II* (Springer-Verlag, Berlin, 1996)
- [DV] D. J. Daley and D. Vere-Jones, *An Introduction to the Theory of Point Processes* (Springer, Berlin, 1988).
- [deS-Z] Ph. de Smedt and V.A. Zagrebnov, Van der Waals Limit of an Interacting Bose Gas in a Weak External Field, *Phys. Rev.* **A35** (1987) 4763–4769.
- [DGPS] F. Dalfovo, S. Giorgini, L. P. Pitaevskii and S. Stringari, Theory of Bose-Einstein condensation in trapped gases, *Rev. Mod. Phys.* **71** (1999) 463–512.
- [EK] N. Eisenbaum and H. Kaspi, On permanental processes, <http://arxiv.org/pdf/math/0610600>.
- [F] W. Freudenberg, Characterization of states of infinite boson systems. II: On the existence of the conditional reduced density matrix. *Commun. Math. Phys.* **137**, 461–472(1991)
- [FF] K.-H. Fichtner and W. Freudenberg, Characterization of states of infinite boson systems. I: On the construction of states of boson systems. *Commun. Math. Phys.* **137**, 315–357(1991)
- [Ka] T. Kato, *Perturbation Theory for Linear Operators* (Springer-Verlag, Heidelberg-Berlin, (1966)).
- [LVZ] J. Lauwers, A. Verbeure and V. A. Zagrebnov, Bose-Einstein Condensation for Homogeneous Interacting Systems with a One-Particle Spectral Gap, *J.Stat.Phys.* **112** (2003) 397–420.
- [LePu] J. Lewis and J. V. Pulè, The Equilibrium States of the Free Boson Gas, *Commun. Math. Phys.* **36** (1974) 1–18 .
- [LSSY] E. H. Lieb, R. Seiringer, J. P. Solovej and J. Yngvason, *The Mathematics of the Bose Gas and its Condensation* (Birkhäuser, Basel (2005)).
- [Ly] E. Lytvynov, Fermion and boson random point processes as particle distributions of infinite free Fermi and Bose gases of finite density, *Rev. Math. Phys.* **14** (2002) 1073–1098.
- [M75] O. Macchi, The coincidence approach to stochastic point processes, *Adv. Appl. Prob.* **7** (1975) 83–122.
- [M77] O. Macchi, The fermion process—a model of stochastic point process with repulsive points, pp.391–398 in *Transactions of the Seventh Prague Conference on Information Theory, Statistical Decision Functions, Random Processes and of*

the Eighth European Meeting of Statisticians (Tech. Univ. Prague, 1974), Vol. A (Reidel, Dordrecht, 1977)

- [MaVe] J. Manuceau, A. Verbeure, Quasi-free states of the CCR-algebra and Bogoliubov transformations, *Commun.Math.Phys.* **9** (1977) 125–131.
- [PeSm] C.J. Pethick and H. Smith, *Bose-Einstein Condensation in Dilute Gases* (Cambridge University Press, Cambridge, (2002)).
- [Pu] J. V. Pulè, The free boson gas in a weak external potential, *J. Math. Phys.* **24** (1983) 138–142.
- [RSIV] M. Reed and B. Simon, *methods of Modern Mathematical Physics*, vol.IV :Analysis of Operatyors (Academic Press, London, (1978)).
- [ShTa] T. Shirai and Y. Takahashi, Random point fields associated with certain Fredholm determinants I: fermion, Poisson and boson point processes, *J. Funct. Anal.* **205** (2003) 414–463.
- [TIa] H. Tamura and K.R. Ito, A Canonical Ensemble Approach to the Fermion/Boson Random Point Processes and its Applications, *Commun. Math. Phys.* **263** (2006) 353–380.
- [TIb] H. Tamura and K.R. Ito, A Random Point Field related to Bose-Einstein Condensation, *J. Funct. Anal.* **243** (2007) 207–231.
- [TIc] H.Tamura and K.R. Ito, Random Point Fields for Para-Particles of Any Order, *J. Math. Phys.* **48** 023301 (2007) (14 pages).
- [TZ] H.Tamura and V.A. Zagrebnov, Boson Random Point Processes in the van der Waals Limit. (in progress)
- [VVZ] L. Vandevenne, A. Verbeure and V. A. Zagrebnov, Equilibrium states for the Bose gas, *J.Math.Phys.* **45** (2004) 1606-1622.
- [V] D. Vere-Jones, A generalization of permanents and determinants, *Linear Algebra Appl.* **111** (1988) 119–124.
- [Z] V.A. Zagrebnov, *Topics in the Theory of Gibbs Semigroups*. Leuven Notes in Mathematical and Theoretical Physics. Vol. 10, *Leuven University Press*, Leuven 2002.
- [ZB] V.A. Zagrebnov and J.-B. Bru, The Bogoliubov model of weakly imperfect Bose gas, *Phys. Rep.* **350** (2001) 291–434 .